

DECOMPOSITION MATRICES AND BLOCKS FOR THE SYMPLECTIC BLOB ALGEBRA OVER THE COMPLEX FIELD

O. H. KING¹, P. P. MARTIN, AND A. E. PARKER

ABSTRACT. We find the blocks for the symplectic blob algebra for all specialisations of the parameters over the complex numbers. We determine Gram determinants for the cell modules. We also show that the algebra is semisimple over the complex numbers unless at least one of the quantisation parameters, or the sum or difference of two of the parameters is integral or q is a root of unity. We also find decomposition numbers in many of the q generic cases.

INTRODUCTION

The *symplectic blob algebra* b_n^x , introduced in [7], is a quotient of the Hecke algebra $H(\tilde{C}_n)$ (Definition 6.1, see for instance [11]). We may define b_n^x using a basis of diagrams which can be thought of loosely as type- $\tilde{C}_n$ Temperley-Lieb diagrams. These are obtained by suitably stacking the ‘decorated’ generators shown in Figure 1 (see [7, §6] or §1 below for details).




FIGURE 1. Generating symplectic blob diagrams.

The algebra b_n^x is defined over any commutative ring k containing a 6-tuple $\boldsymbol{\delta} = (\delta, \delta_L, \delta_R, \kappa_L, \kappa_R, \kappa_{LR})$ of ‘parameters’. In this paper we study representation theory. As such our aim is primarily the Artinian cases, and indeed the cases where k is an algebraically closed field. Thus for each such k there is an algebra b_n^x for each point in k^6 . One knows [7] (and see §7) that the non-semisimple cases lie on certain algebraic sets. The generic semisimple cases are well-understood [7], so it is the points on the algebraic sets that are of interest.

It turns out that the dependence of representation theory on position in the variety is more easily described in terms of alternative variety-specific parameterisations. In these the non-semisimple sub-varieties correspond to integrality of some or all of the new parameters, as we shall see in §8 (see also [17, 5]). In particular we introduce parameter q and set $\delta = q + q^{-1}$.

¹Corresponding author

In [7] various general properties of the algebra b_n^x are established. For instance a cellular basis is constructed; its generic semisimple structure over $\mathbb{C}$ is determined; and it is shown to be quasi-hereditary on an open subvariety of the non-semisimple variety. Full tilting modules are constructed in [19]. An efficient presentation is found in [8]; and in [5] a closely related algebra is studied, leading to useful alternative bases for certain cell modules, which are crucial to our calculation of the action of a certain central element.

It follows from comparison with the ordinary blob algebra case [4] that the programme of study of the non-generic non-semisimple representation theory of b_n^x is a considerably harder challenge. As in [4], a key component is to construct ‘enough’ standard module morphisms and these were constructed in [9]. This paper, using the morphisms and also using [5], investigates the sufficiency of this set.

Quite generally, if there is a non-zero homomorphism between two standard modules, then the two modules must belong to the same block. Indeed, determination of *all* homomorphisms between standard modules in a quasi-hereditary structure allows a complete description of the blocks (see the appendix). Our main block result in this paper, contained in §10–11, gives the blocks in the *subcritical* case, namely in characteristic zero, q not a root of unity.

The homomorphisms found in [9] are not shown to be a complete set, so only give a lower bound on the size of blocks. However these results combined with a result about the action of certain central elements on the standard modules allow us to obtain an *upper* bound on the size of blocks. The homomorphisms (along with some restriction results to the blob algebra) then allow a complete characterisation of the blocks.

The paper is structured as follows. We give a brief review of notation in section 1, and of the construction of cell modules in §2. In §3 we discuss the role of the ground ring. In §4 we review the de Gier–Nichols path basis of cell modules. The first main theorems are in §6, which gives conditions for two cell modules to be in the same block. In §7 we compute Gram determinants, and in §9–10 the main theorems on decomposition matrices and geometric characterisation of blocks are given.


1. NOTATION AND PRELIMINARY DEFINITIONS

Let k be a field and

$$\boldsymbol{\delta} = (\delta, \delta_L, \delta_R, \kappa_L, \kappa_R, \kappa_{LR}) \in k^6. \quad (1)$$

Let $\mathbb{N}_0$ denote the natural numbers including 0. Let $n, m \in \mathbb{N}_0$.

The set $B_n^{x'}$ of *left-right blob pseudo-diagrams* [9] may be defined as follows. Consider the set of decorated Temperley–Lieb diagrams on n strings in Figure 1 (as usual for Temperley–Lieb diagrams, isotopic pictures are identified). Then $B_n^{x'}$ is the set of pictures (up to isotopy) obtained by stacking such pictures. Write $d|d'$ for diagram d stacked over diagram d' .

TABLE 1. Some features appearing in diagrams in the set $B_n^{x'}$.

δ	δ_L	δ_R	κ_L	κ_R	κ_{LR}	κ_{LR}	κ_{LR}

TABLE 2. Table encoding straightening relations for b_n^x .

Let B_n^x denote the subset of $B_n^{x'}$ excluding diagrams with features as in Table 1. Given $d \in B_n^{x'}$, an element $f(d)$ of kB_n^x is obtained by applying the straightening relations encoded in Table 2 (the feature on the top is replaced by the given scalar multiple of the feature beneath) and the “topological relation”:

$$\kappa_{LR} \left(\begin{array}{c} \text{Diagram with shaded regions C, A, B, D and a vertical line with a black dot} \end{array} \right) = \begin{array}{c} \text{Diagram with shaded regions C, A, B, D and a vertical line with a white dot} \end{array} \quad (2)$$

(where each labelled shaded area is a subdiagram without propagating lines) until such operations are exhausted. It is shown in [9] that $f(d)$ does not depend on the details. Thus we have in particular a well-defined map $B_n^x \times B_n^x \rightarrow kB_n^x$ given by

$$(d, d') \mapsto f(d|d'). \quad (3)$$

Definition 1.1. Fix k and $\delta \in k^6$. Then the *symplectic blob algebra* b_n^x is the k -algebra with basis B_n^x , and multiplication as in (3).

For example, consider the poset $(\Lambda_n, \prec)$ given in Figure 2(a) and the elements d_l ($l \in \Lambda_n$) of b_n^x as indicated in Figure 3, where in particular

$$d_0 = \begin{array}{c} \text{Diagram with } n \text{ black dots on a horizontal line} \end{array} \quad \text{if } n \text{ is even,} \quad d_0 = \begin{array}{c} \text{Diagram with } n \text{ black dots on a horizontal line and a vertical line on the right} \end{array} \quad \text{if } n \text{ is odd.}$$

Verification of the following is a simple exercise in b_n^x arithmetic.

Proposition 1.2 ([7]). *The ideals of b_n^x generated by the elements d_l in Figure 3 include according to the poset structure indicated by the Hasse diagram in Figure 2.* $\square$

Let $I_n(0)$ denote the ideal generated by d_0 . Define b'_n as the quotient algebra by this ideal.




FIGURE 2. (a) Cell-module weight-label poset. Here 0 is the maximal element and $-n$ is the minimal element. (b) Cell-module weight label poset in DN-labelling (even $n = 2m$ case), see §2.1.

2. REVIEW OF CONSTRUCTION OF b_n^x CELL MODULES

Consider the poset $(\Lambda_n, \prec)$ given in Figure 2. A set $\{S_n(l)\}_{l \in \Lambda_n}$ of b_n^x -modules is constructed over arbitrary k in [7]. In this section we review the construction. One should start by thinking of k not as a field but rather as the commutative ring $\mathbb{Z}[\delta]$ here. Then one can pass to any case by base change. These modules pass to simple modules in the semisimple cases (see [7]), so they can be thought of as the integral forms of the ‘ordinary’ irreducibles in a Brauer-modular system [3, 2]. (Although our setup requires careful preparation to be properly modular, cf. [2] — we will not need to develop the full machinery here.)

The left b_n^x -module $S_n(l)$ has a basis of half-diagrams constructed similarly to the blob algebra case (cf. [4, p. 593], [7, Section 8]). See Figure 4 for an example. Note that by (3) the left action corresponds to stacking a diagram on top of the basis element.

Consider $l \in \Lambda_n$. To construct a basis $\beta_n(l)$ for b_n^x -module $S_n(l)$ in general we proceed as follows. Consider the subset of B_n^x of diagrams with $|l|$ undecorated propagating lines. If l is positive, then further restrict to diagrams with a left blob on the first propagating line. Otherwise, if l is negative, then there must be no such blob. Now pick any one of the remaining diagrams d , and take the subset of diagrams agreeing with d in the lower half. Finally, as the lower half is the same in all diagrams, and does not affect multiplication, we omit it. (As another example, half-diagram bases for the cell modules for low rank b_n^x are listed in [7, Figure 3]. There cut lines are used in place





FIGURE 3. Representative diagrams in the cell ideal poset.

FIGURE 4. Diagram basis of cell module $S_5(1) = W_{-, -}^{(5,2)}$.

of blobs.) The algebra action is by diagram stacking, except that diagrams arising that lie outside the basis (necessarily with higher weight in the sense of Figure 2) are zero. Recall:

Proposition 2.1 ([7]). *The algebra b_n^x is a cellular algebra (in the sense of [6]). The modules $\{S_n(l)\}_{l \in \Lambda_n}$ are the cell modules. The labelling poset Λ_n for the cell modules is as in Figure 2.*

When all parameters are invertible, Λ_n also labels the simple modules, in which case the algebra is also quasi-hereditary with the above poset and the cell modules are standard modules.

2.1. On standard and De Gier–Nichols weight labelling. In [5] there is a useful reformulation of $(\Lambda_n, \prec)$ as follows. The basis B_n^x is equivalent to a basis of affine-symmetric TL (ASTL) diagrams (see [7] for the equivalence). In an ASTL diagram “blobs” are indicated by paired lines that touch the left (for a left blob) or the right (a right blob) side of a diagram. A corresponding half-diagram can in principle have any number of lines touching the left or right side, but the parity of each number is preserved in the (ASTL version of the) basis of a cell module. Thus for $\varepsilon_i \in \{\pm 1\}$ the module $W_{\varepsilon_1, \varepsilon_2}^{n, m}$ is the cell module with ASTL half-diagram basis with ε_1 parity on the left side, ε_2 parity on the right side and $m + \frac{1}{2}(\varepsilon_1 + \varepsilon_2)$ propagating lines (here $\varepsilon_i = +1$ for even, written as $+$; and $\varepsilon_i = -1$ for odd, written as $-$). We write Λ_n^+ for the new labelling scheme — see Figure 2(b).

The correspondence (in both directions) is given as follows:

$$S_n(l) = \begin{cases} W_{-\text{sgn } l, \text{sgn } l}^{(n, |l|)} & \text{if } n \text{ and } l \text{ have opposite parity, } l \neq 0, \\ W_{-\text{sgn } l, -\text{sgn } l}^{(n, |l+1|)} & \text{if } n \text{ and } l \text{ have the same parity, } l \neq 0, \\ W^n(b) & \text{if } l = 0 \end{cases}$$

$$W_{\varepsilon_1, \varepsilon_2}^{(n, l)} = S_n(-\varepsilon_1(l + \frac{1}{2}(\varepsilon_1 + \varepsilon_2))) \quad (4)$$

where $\text{sgn } l$ is the sign of l .

Remark: The argument b used for the cell module with no propagating lines indicates that this module depends on a parameter b . This is essentially the same as κ_{LR} (see §3.2).

3. ON δ PARAMETER CONVENTIONS AND REPARAMETERISATION

3.1. Ground ring arithmetic. In the modular system [3] one works largely in the integral ground ring, passing to a specific modular case (to address specific Artinian representation theory) as late as possible. However for reasons of arithmetic manipulation it may be expedient to perform

	generator scaling			parameter scaling / reparameterisation					
label	$e \mapsto$	$e_i \mapsto$	$f \mapsto$	$\delta \mapsto$	$\delta_L \mapsto$	$\delta_R \mapsto$	$\kappa_L \mapsto$	$\kappa_R \mapsto$	$\kappa_{LR} \mapsto$
1	$\frac{e}{\kappa_L}$	e_i	$\frac{f}{\kappa_R}$	δ	$\frac{\delta_L}{\kappa_L}$	$\frac{\delta_R}{\kappa_R}$	1	1	$\frac{\kappa_{LR}}{\kappa_L \kappa_R}$
2	$-e$	$-e_i$	$-f$	$-\delta$	$-\delta_L$	$-\delta_R$	κ_L	κ_R	κ_{LR}
DN				$[2]$	$\frac{[\omega_1]}{[\omega_1+1]}$	$\frac{[\omega_2]}{[\omega_2+1]}$	1	1	b
GMP1				$[2]$	$[w_1]$	$[w_2]$	$[w_1 + 1]$	$[w_2 + 1]$	κ_{LR}
GMP2				$-[2]$	$-[w_1]$	$-[w_2]$	$[w_1 + 1]$	$[w_2 + 1]$	κ_{LR}

TABLE 3. Alternative parameterisations for b_n^x .

computations as if in a different ground ring. This looks like base-change away from the generality of the integral ring. But provided the change is arithmetically reversible back to the integral ring, it is not restrictive.

An example is as follows. The substitution homomorphism $\mathbb{Z}[\delta] \rightarrow \mathbb{Z}[q, q^{-1}]$ given by $\delta \mapsto q + q^{-1}$ is not an isomorphism. However it is an injection, so the map can be inverted on any element of the image. Thus one can do arithmetic on elements of $\mathbb{Z}[\delta]$ working in the image, and then recover identities that hold in $\mathbb{Z}[\delta]$.

In the example, a merit of the substitution is if one works with elements of $\mathbb{Z}[\delta]$ satisfying the recursion $p_n = [2]p_{n-1} - p_{n-2}$, with $p_0 = 0$ and $p_1 = 1$ (for example certain Gram determinants of the Temperley–Lieb algebra satisfy this recursion [17]). This is the Tchebychev recursion [16, §6.3.3]. The complex roots of p_n are the so-called Beraha numbers [1] — but factorisation is not obvious. However working in the image these elements take the simple form $p_n = [n]$, where

$$[m] := q^{-m+1} + q^{-m+3} + \cdots + q^{m-3} + q^{m-1}.$$

This formulation has manifest factorisation properties. In particular $[n] = 0$ requires q to be a root of unity.

3.2. Parameterisation by exponents w_1, w_2 . In order to determine the representation theory of $b_n^x(\delta)$ it is useful to reparameterise as discussed in [9, §2]. We recall the key parameterisations in Table 3. Generator scaling “1” in Table 3 induces an isomorphism with the algebra with parameters rescaled as shown, reducing from 6 parameters to 4 [8]. “GMP1”, “GMP2” and “DN” reparameterise with parameters q, w_1, w_2 (cf. [1, 14, 15, 20, 17]). DN is the parameter choice of de Gier–Nichols in [5]. GMP1 and GMP2 are the parameter choices that were most useful for [9]. GMP1 and GMP2 can be converted from one to another by taking the isomorphic algebra with generators multiplied by -1 , i.e. using “2” to rescale. GMP2 turns out to be the most convenient for presenting the results about general families of homomorphisms in [9]. Then “1” converts from DN to GMP1 and then to GMP2 via “2”.

De Gier–Nichols further reparameterise b in terms of a new parameter θ :

$$b = \begin{cases} \left[\frac{w_1 + w_2 + \theta + 1}{2} \right] \left[\frac{w_1 + w_2 - \theta + 1}{2} \right] & \text{if } n \text{ even} \\ - \left[\frac{w_1 - w_2 + \theta}{2} \right] \left[\frac{w_1 - w_2 - \theta}{2} \right] & \text{if } n \text{ odd.} \end{cases} \quad (5)$$

4. BASES OF THE b_n^x -MODULE $W^{(n)}(b)$

Definition 4.1. For $n \in \mathbb{N}$ a *path* p is an element of the set

$$\mathcal{P}_n = \{ p = (h_0, h_1, \dots, h_n) \mid h_0 = 0; \text{ and } |h_{i+1} - h_i| = 1 \text{ for } 0 \leq i \leq n-1 \}$$

In particular, define the *fundamental path* $p_0 = (0, -1, 0, -1, 0, \dots)$.




FIGURE 5. The tiled lattice and the fundamental path p_0 .

We can draw paths on a tiled square lattice as in Figure 5. Each path p may be partitioned according to the points at which it agrees with p_0 . There are in particular parts of p that are above or below p_0 . Each of these parts defines an ‘envelope’ between the two paths. We can move from p_0 to p through a sequence of intermediate paths p_i by ‘adding tiles’ (or half tiles) within each envelope. In particular note that if $p_i \neq p$ then there is always a lowest numbered position at which a tile can be added. Define $\mathcal{P}(p)$ as the ordered set passing from p_0 to p in this way.

Define E'_n in b_n^x by $E'_n = d_0$. That is:

$$E'_n = \begin{array}{|c|} \hline \bullet \quad \cdots \quad \bullet \\ \hline \bullet \quad \cdots \quad \bullet \\ \hline \end{array} \quad \text{if } n \text{ is even,} \quad E'_n = \begin{array}{|c|} \hline \bullet \quad \cdots \quad \bullet \\ \hline \bullet \quad \cdots \quad \bullet \quad \bullet \\ \hline \end{array} \quad \text{if } n \text{ is odd.}$$

Note from (4) that

$$W^{(n)}(b) = b_n^x E'_n.$$

Define a subset $\pi_n = \{w_p | p \in \mathcal{P}_n\}$ of $W^{(n)}(b)$ as follows. To a path p we associate an element w_p defined recursively through $\mathcal{P}(p)$: firstly $w_{p_0} = E'_n$; then $w_{p_{j+1}} = e_i w_{p_j}$ if p_{j+1} obtained from p_j by adding a tile in position i .

Theorem 4.2. *The subset π_n is the diagram basis for $W^{(n)}(b)$.* $\square$

4.1. Path basis for $W^{(n)}(b)$ for generic δ .

Here we will use a notion of *generic* $\delta \in k^6$ [10]. A point is generic if it lies in the (Zariski) open subset excluding a certain variety (in our case the variety given by the collection of denominators in a construction below — see (6)). The utility is that every δ in $\mathbb{C}^6$ is the limit of a set of generic points, so that certain identities $f(\delta) = 0$ that hold generically will hold at every point where f makes sense.

We define a formal subset of the b_n^x -module $W^{(n)}(b) = b_n^x E'_n$ for generic δ . To a path p we associate an element v_p , defined recursively through $\mathcal{P}(p)$ as follows:

$$\begin{aligned} v_{p_0} &= E'_n, \\ v_{p'} &= X_i v_p \text{ if } p' \text{ is obtained from } p \text{ by adding a tile at position } i, \end{aligned}$$

where X_i is one of the following operators:

- $X_i = e_i - r(w_1 - h_{i-1})1$ if a full tile is added from above;
- $X_i = e_n - k(w_1 - h_{n-1})1$ if a half tile is added from above at the right boundary;
- $X_i = e_i - r(-w_1 + h_{i-1})1$ if a full tile is added from below;
- $X_i = e_n - k(-w_1 + h_{n-1})1$ if a half tile is added from below at the right boundary,

where

$$r(u) = \frac{[u+1]}{[u]}, \quad \text{and } k(u) = -\frac{[(u-w_2+\theta)/2][(u-w_2-\theta)/2]}{[u][w_2+1]}. \quad (6)$$

Define

$$\Pi_n = \{v_p | p \in \mathcal{P}_n\}$$

Comparing the constructions we have immediately from Theorem 4.2:

Theorem 4.3. *When defined, the set Π_n can be obtained from π_n by an upper-unitriangular transformation; and hence is a basis for $W^{(n)}(b)$.* $\square$

Theorem 4.4. *In general there are other ways of adding tiles to pass from p_0 to each p (cf. the ordered sequence $\mathcal{P}(p)$). The construction does not depend on the choice of route.*

Proof. For each p note that if there are two routes to p from some lower path then the different sequences of multiplications involve pairwise commuting factors. $\square$

Remark. If the scalar term is omitted in X_i , this construction builds the diagram basis, up to the DN rescaling factors. We shall later keep track of these scalars explicitly, and hence recover

‘integral-valued’ Gram matrices from certain nominal Gram matrix calculations. (It is interesting to contrast this with the path basis for Temperley–Lieb in [16]. There the orthogonal basis is orthonormal, so the nominal Gram matrix is the identity matrix, and one *only* has to work out the basis scaling factor.)

Theorem 4.5 ([5, Theorem 5.9]). *Let $p = (h_0, h_1, \dots, h_n) \in \mathcal{P}_n$. Then the generators $e = e_0, e_1, \dots, e_n = f$ have the following action on v_p :*

- *Each v_p is an eigenvector for the left blob generator e_0 :*
 - (1) *If $h_1 = -1$ then $e_0 v_p = \frac{[w_1]}{[w_1+1]} v_p$.*
 - (2) *If $h_1 = 1$ then $e_0 v_p = 0$.*
- *The action of e_i ($1 \leq i \leq n-1$) on v_p is zero if p has positive or negative slope at position i , i.e. $|h_{i-1} - h_{i+1}| = 2$. If this is not the case, then let p' be the path obtained by adding a tile to p at position i . Then e_i acts on the pair $\{v_p, v_{p'}\}$ in the following way:*

- (1) *If $h_{i-1} \geq 0$ then*

$$\begin{aligned} e_i v_p &= v_{p'} + r(w_1 - h_{i-1}) v_p \\ e_i v_{p'} &= r(-w_1 + h_{i-1}) v_{p'} + r(-w_1 + h_{i-1}) r(w_1 - h_{i-1}) v_p. \end{aligned}$$

- (2) *If $h_{i-1} < 0$ then*

$$\begin{aligned} e_i v_p &= v_{p'} + r(-w_1 + h_{i-1}) v_p \\ e_i v_{p'} &= r(w_1 - h_{i-1}) v_{p'} + r(-w_1 + h_{i-1}) r(w_1 - h_{i-1}) v_p. \end{aligned}$$

- *Let p' be the path obtained by adding a half tile to p at the right boundary. Then e_n acts on the pair $\{v_p, v_{p'}\}$ in the following way:*

- (1) *If $h_{n-1} \geq 0$ then*

$$\begin{aligned} e_n v_p &= v_{p'} + k(w_1 - h_{n-1}) v_p \\ e_n v_{p'} &= k(-w_1 + h_{n-1}) v_{p'} + k(-w_1 + h_{n-1}) k(w_1 - h_{n-1}) v_p. \end{aligned}$$

- (2) *If $h_{n-1} < 0$ then*

$$\begin{aligned} e_n v_p &= v_{p'} + k(-w_1 + h_{n-1}) v_p \\ e_n v_{p'} &= k(w_1 - h_{n-1}) v_{p'} + k(-w_1 + h_{n-1}) k(w_1 - h_{n-1}) v_p. \end{aligned}$$

5. RESTRICTING STANDARD MODULES TO THE BLOB ALGEBRA

The (left) blob algebra b_n is the subalgebra of b_n^x generated by $\{e_0, e_1, \dots, e_{n-1}\}$ [17]. The generators $\{e_1, \dots, e_{n-1}, e_n\}$ generate another copy of b_n which we will call the *right blob algebra*.

In [7, §8] the restriction to b_n is used to determine the dimensions of the standard modules, $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$. There it is shown that each restricted $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ is filtered by standard b_n -modules (as

	left blob module		right blob module	
	$\varepsilon_2 = 1$	$\varepsilon_2 = -1$	$\varepsilon_2 = 1$	$\varepsilon_2 = -1$
$\varepsilon_1 = 1$	$W_n(n)$	$W_n(n)$	$W_n(n)$	$W_{-n}(n)$
	$\vdots$	$\vdots$	$\vdots$	$\vdots$
	$W_{m+3}(n)$	$W_{m+3}(n)$	$W_{m+3}(n)$	$W_{-(m+3)}(n)$
	$W_{m+1}(n)$	$W_{m+1}(n)$	$W_{m+1}(n)$	$W_{-(m+1)}(n)$
$\varepsilon_1 = -1$	$W_{-n}(n)$	$W_{-n}(n)$	$W_n(n)$	$W_{-n}(n)$
	$\vdots$	$\vdots$	$\vdots$	$\vdots$
	$W_{-(m+3)}(n)$	$W_{-(m+3)}(n)$	$W_{m+3}(n)$	$W_{-(m+3)}(n)$
	$W_{-(m+1)}(n)$	$W_{-(m+1)}(n)$	$W_{m+1}(n)$	$W_{-(m+1)}(n)$

TABLE 4. The standard content of $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ as a left (resp. right) blob module.

defined in [17] — the construction is analogous to §2). We follow the notation of [4] and use $W_{\pm t}(n)$ for the standard b_n -modules. Recall that $W_t(n)$ is the standard blob module with half diagram basis that has n northern nodes and t (undecorated) propagating lines. $W_{-t}(n)$ is the standard blob module with half diagram basis that has n northern nodes and $t - 1$ (undecorated) propagating lines and one decorated propagating line.

The restriction will again be useful here. Any b_n^x -homomorphism is also a left (right) blob homomorphism upon restriction, and thus must respect any left (right) blob structure.

Let d be a half diagram that generates some $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$, as in §2. We define $\text{ur}_1(d)$ to be the number of lines crossing the 1-wall (in the sense of [7], i.e. the right wall), not counting any lines that are part of non-contractible loops. We similarly define $\text{ur}_0(d)$ as the number of lines crossing the left wall.

When we restrict $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ to the left blob algebra then it is filtered by ur_1 and each section is isomorphic to a standard blob module. A similar situation occurs when we restrict to the right blob algebra. We have the following.

Proposition 5.1 ([7]). *The b_n -standard content of $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ is as in Table 4.* □

When the left (or right) blob algebra is semi-simple, then every standard module is simple.

6. A NECESSARY BLOCK CONDITION

In this section we recall a central element Z_n (see (19) below) of b_n^x . We prove Conjecture 6.5 from [5] and deduce the action of Z_n on cell modules. We use this to investigate the block structure.

6.1. The central element Z_n . We shall need a surjection from $H(\tilde{C}_n)$ to b_n^x as in [7, Proposition 6.3.2]. Further details can be found in [5, §2] (caveat: there are typos in [5]; cf. e.g. [14]).

Definition 6.1. Let q, Q_1 and Q_2 be indeterminates. The Hecke algebra $H(\tilde{C}_n)$ of type $\tilde{C}_n$ over $\mathbb{Z}[q^{\pm 1}, Q_1^{\pm 1}, Q_2^{\pm 1}]$ is the associative algebra with generators $g_0, g_1, \dots, g_n$ and relations:

$$g_i g_{i+1} g_i = g_{i+1} g_i g_{i+1}, \quad 1 \leq i \leq n-2, \quad (7)$$

$$g_0 g_1 g_0 g_1 = g_1 g_0 g_1 g_0, \quad (8)$$

$$g_{n-1} g_n g_{n-1} g_n = g_n g_{n-1} g_n g_{n-1}, \quad (9)$$

$$g_i g_j = g_j g_i, \quad |i - j| > 1, \quad (10)$$

$$(g_i - q)(g_i + q^{-1}) = 0, \quad 1 \leq i \leq n-2, \quad (11)$$

$$(g_0 - Q_1)(g_0 + Q_1^{-1}) = 0, \quad (12)$$

$$(g_n - Q_2)(g_n + Q_2^{-1}) = 0. \quad (13)$$

For suitable base change and choices of the parameters we have successive quotients:

$$H(\tilde{C}_n) \rightarrow 2\text{BTL} \rightarrow b_n^x \rightarrow \frac{b_n^x}{I_n(0)} \quad (14)$$

where 2BTL is defined in [8].

The algebra b_n^x is defined over a ring k with parameters $\delta = (\delta, \delta_L, \delta_R, \kappa_L, \kappa_R, \kappa_{LR}) \in k^6$. For any three units in k we can view k as a $\mathbb{Z}[q^{\pm 1}, Q_1^{\pm 1}, Q_2^{\pm 1}]$ -algebra by making q, Q_1 and Q_2 act as these units. For each such triple we understand $H(\tilde{C}_n)$ as a k -algebra by base change.

Note that we are using the Saleur normalisation [17] for generators.

Proposition 6.2. *By abuse of notation let us write q, Q_1, Q_2 for the actions of these three scalars in k defining $H(\tilde{C}_n)$ as a k -algebra as described above. If they satisfy*

$$\delta = [2], \quad (qQ_1 - q^{-1}Q_1^{-1})\delta_L = Q_1 - Q_1^{-1}, \quad (qQ_2 - q^{-1}Q_2^{-1})\delta_R = Q_2 - Q_2^{-1}. \quad (15)$$

then there is a surjective k -algebra homomorphism $\pi : H(\tilde{C}_n) \rightarrow b_n^x$, given by

$$\pi(g_i^{\pm 1}) = e_i - q^{\mp 1}, \quad (16)$$

$$\pi(g_0^{\pm 1}) = Q_1^{\pm 1} - (q^{\pm 1}Q_1^{\pm 1} - q^{\mp 1}Q_1^{\mp 1})e_0, \quad (17)$$

$$\pi(g_n^{\pm 1}) = Q_2^{\pm 1} - (q^{\pm 1}Q_2^{\pm 1} - q^{\mp 1}Q_2^{\mp 1})e_n. \quad (18)$$

(Note that there is no dependence on κ_{LR} .)

Proof. (Outline) Consider (11):

$$\pi(g_i)\pi(g_i) = (e_i - q^{-1})(e_i - q^{-1}) = \delta e_i - 2q^{-1}e_i + q^{-2} = (q - q^{-1})e_i + q^{-2}$$

$$\pi(g_i^2) \stackrel{(11)}{=} \pi((q - q^{-1})g_i + 1) = (q - q^{-1})(e_i - q^{-1}) + 1 = (q - q^{-1})e_i + q^{-2}$$

Alternatively here note that by (11) g_i has eigenvalues q and $-q^{-1}$; while e_i has eigenvalues $\delta, 0$. Then $e_i - q^{-1}$ has eigenvalues $q, -q^{-1}$, by (15), as required. Similarly by (12) g_0 has eigenvalues $Q_1, -Q_1^{-1}$; while e_0 has eigenvalues $\delta_L, 0$. Then $Q_1 - (qQ_1 - q^{-1}Q_1^{-1})e_0$ has eigenvalues $Q_1, -Q_1^{-1}$ as required provided that (15) holds. The verification for g_n is directly analogous. $\square$

The homomorphism π allows elements of $H(\tilde{C}_n)$ to act on b_n^x . In particular,

Definition 6.3 ([5, Definition 2.8]). The ‘Murphy elements’ for $H(\tilde{C}_n)$ are:

$$J_0 = g_1^{-1}g_2^{-1} \cdots g_{n-1}^{-1}g_ng_{n-1} \cdots g_2g_1g_0$$

$$J_i = g_iJ_{i-1}g_i, \quad 1 \leq i \leq n-1.$$

Proposition 6.4 ([5, Proposition 2.10]). *The Murphy elements J_i are pairwise commuting and obey the following relations:*

$$[g_0, J_j] = 0, \quad j \neq 0,$$

$$[g_i, J_j] = 0, \quad 1 \leq i \leq n-1, j \neq i-1, i,$$

$$[g_i, J_{i-1}J_i] = 0, \quad 1 \leq i \leq n-1,$$

$$[g_i, J_{i-1} + J_i] = 0, \quad 1 \leq i \leq n-1,$$

$$[g_0, J_0 + J_0^{-1}] = 0.$$

In particular, the completely symmetric polynomials in J_i, J_i^{-1} ($0 \leq i \leq n-1$) are central in $H(\tilde{C}_n)$.

We hence let Z_n be the central element

$$Z_n = \sum_{i=0}^{n-1} (J_i + J_i^{-1}). \tag{19}$$

6.2. Aside on substitutions. For $w_1, w_2 \in \mathbb{Z}$ we set

$$Q_1 = q^{w_1} \text{ and } Q_2 = q^{w_2},$$

so that (15) becomes

$$\delta = [2], \quad \delta_L = \frac{[w_1]}{[w_1 + 1]}, \quad \delta_R = \frac{[w_2]}{[w_2 + 1]}.$$

We can interpret $[w_1 + a]$ ($a \in \mathbb{Z}$) in the following way:

$$[w_1 + a] = \frac{q^{w_1+a} - q^{-w_1-a}}{q - q^{-1}} = \frac{Q_1q^a - Q_1^{-1}q^{-a}}{q - q^{-1}}.$$

Similarly we have

$$[w_2 + a] = \frac{Q_2 q^a - Q_2^{-1} q^{-a}}{q - q^{-1}} \quad \text{and} \quad [w_1 + w_2 + a] = \frac{Q_1 Q_2 q^a - Q_1^{-1} Q_2^{-1} q^{-a}}{q - q^{-1}}.$$

6.3. The Z_n -action theorem. The following lemma is mostly a restatement of [5, Proposition 5.19]. However we have also included the labels of the irreducible modules.

Lemma 6.5. *The generic b_n -module with basis $\{v_p \mid p \text{ of fixed final height } h_n\}$ in $W^{(n)}(b)$ is isomorphic to the generic irreducible b_n -module $W_{h_n}(n)$.*

Proof. Note first that this is indeed a module for the blob algebra, as the only elements of the symplectic blob algebra that change the final height of a path involve the generator e_n , which is not present when we consider the restricted action.

Now by [5, Proposition 5.19] these modules are the generic irreducibles for the blob algebra, so it suffices to show that the labelling matches up.

First consider the case when n is even. From [18, (3.2)] we have a maximal heredity chain of idempotents

$$\left(\frac{1}{\delta^{n/2}} e_1 e_3 \dots e_{n-1}, \frac{1}{\delta_L \delta^{n/2-1}} e_0 e_3 e_5 \dots e_{n-1}, \dots, \frac{1}{\delta} e_{n-1}, \frac{1}{\delta_L} e_0, 1 \right) \quad (20)$$

corresponding to the standard modules $W_0(n), W_{-2}(n), \dots, W_{n-2}(n), W_{-n}(n), W_n(n)$ respectively. We must therefore show that the module with basis $\{v_p \mid p \text{ with fixed final height}\}$ is associated to the correct heredity idempotent. Suppose first that the final height is $h_n = 0$, then this module contains the element v_{p_0} , where p_0 is the fundamental path. By Theorem 4.5 none of the idempotents in (20) kill v_{p_0} , therefore this module must be isomorphic to $W_0(n)$.

If now the final height is $h_n > 0$, then all paths must either have a slope at at least h_n points, or start with $h_1 = 1$ and have a slope at at least $h_n - 1$ points. Since the e_i in the heredity idempotents commute, we therefore see that any idempotent containing a product of at least $(n - h_n)/2 + 1$ of the e_i will kill the basis elements obtained from these paths, but those containing $(n - h_n)/2$ will not. Therefore the first heredity idempotent that does not annihilate this module is

$$\frac{1}{\delta^{(n-h_n)/2}} e_{h_n+1} \dots e_{n-1},$$

which corresponds to the left blob module $W_{h_n}(n)$.

If the final height is $h_n < 0$, then again all paths must have a slope at at least $|h_n|$ points, or start with $h_1 = -1$ and have a slope at at least $h_n - 1$ points. In this case, any idempotent containing a product of at least $(n + h_n)/2 + 1$ of the e_i for $i \neq 0$ will kill the basis elements obtained from these paths, but those containing a product of $(n + h_n)/2$ of the e_i ($i \neq 0$) and e_0 will not. Therefore the first heredity idempotent that does not annihilate this module is

$$\frac{1}{\delta_L \delta^{(n+h_n)/2}} e_0 e_{-h_n+1} \dots e_{n-1},$$

which corresponds to the left blob module $W_{h_n}(n)$.

The proof for n odd is similar. $\square$

We now use the path basis to determine submodules of $W^{(n)}(b)$ for specific parameter choices.

Proposition 6.6 ([5, Proposition 6.3]). *Fix $m, n \in \mathbb{N}_0$ and $\varepsilon_2 \in \{\pm 1\}$. Fix $\delta \in k^6$ except for κ_{LR} , generic, but so that w_1, w_2 are defined.*

- (i) *Choose θ so that $[(-m + w_1 + \varepsilon_2 w_2 \pm \theta)/2] = 0$. Then the b_n^x -module $W^{(n)}(b)$ has a submodule $V_{+, \varepsilon_2}^{(n, m)}$ with basis $\pi_n^+(m) = \{v_p \mid p = (h_0, h_1, \dots, h_n) \text{ with } h_n \geq m + 1\}$.*
- (ii) *Choose θ so that $[(-m - w_1 + \varepsilon_2 w_2 \pm \theta)/2] = 0$. Then $W^{(n)}(b)$ has a submodule $V_{-, \varepsilon_2}^{(n, m)}$ with basis $\pi_n^-(m) = \{v_p \mid p = (h_0, h_1, \dots, h_n) \text{ with } h_n \leq -m - 1\}$.*

This statement is slightly modified from [5]. The key point is that $k(\pm(w_1 - h_{N-1}))$ is zero, and this is equivalent to requiring $[(-m \pm w_1 \pm w_2 \pm \theta)/2] = 0$ for appropriate signs.

Theorem 6.7 ([5, Theorem 6.4]). *Fix $\delta \in k^6$ except for κ_{LR} , generic, but so that w_1, w_2 are defined. Fix $\frac{n, m}{\varepsilon_1, \varepsilon_2} \in \Lambda_n^+$. Let $\theta = -m + \varepsilon_1 w_1 + \varepsilon_2 w_2$. The action of the central element Z_n as defined in (19) on $V_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ as defined in Proposition 6.6 is given by*

$$Z_n V_{\varepsilon_1, \varepsilon_2}^{(n, m)} = \alpha_{\varepsilon_1, \varepsilon_2}^{(n, m)} V_{\varepsilon_1, \varepsilon_2}^{(n, m)}$$

where

$$\alpha_{\varepsilon_1, \varepsilon_2}^{(n, m)} = [n] \frac{[2(-m + \varepsilon_1 w_1 + \varepsilon_2 w_2)]}{[-m + \varepsilon_1 w_1 + \varepsilon_2 w_2]}.$$


Theorem 6.8. *Let $\theta = -m + \varepsilon_1 w_1 + \varepsilon_2 w_2$. Then the generic b_n^x -module $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ is isomorphic to the submodule $V_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ of $W^{(n)}(b)$.*

Proof. Consider first the case $\varepsilon_1 = \varepsilon_2 = 1$. Let $d_{m+1} \in W_{+, +}^{(n, m)}$ be as in (21).

$$d_{m+1} = \begin{array}{|c|c|c|c|c|} \hline \bullet & \bullet & & & \\ \hline \cdots & & & & \\ \hline \bullet & \bullet & & & \\ \hline \end{array} \quad \underbrace{\hspace{1.5cm}}_{m+1} \quad (21)$$

Note d_{m+1} generates $W_{+, +}^{(n, m)}$. Let p_{m+1} be the path of height $h_n = m + 1$ which can be obtained from the fundamental path p_0 by adding the minimal number of tiles. For instance, if $n = 6$ and

$h_n = 2$ then p_2 is the path below.



We claim that we have a morphism $\phi : W_{+,+}^{(n,m)} \longrightarrow V_{+,+}^{(n,m)}$ that is generated by sending $d_{m+1} \mapsto v_{p_{m+1}}$. We first show that the subset of the symplectic blob generators that annihilate d_{m+1} also annihilate $v_{p_{m+1}}$. Indeed we must consider the elements $e_{n-m-1}, \dots, e_n$. But at the steps corresponding to these indices (except n) the path p_{m+1} has a slope, so by Theorem 4.5 the element $v_{p_{m+1}}$ is annihilated by these generators. For the action of e_n , note also from Theorem 4.5 that our choice of θ causes e_n to kill $v_{p_{m+1}}$ also. Therefore ϕ is a homomorphism.

We now show by induction that the homomorphism ϕ is surjective. Define a partial order on the set of paths of final height at least $m+1$ by $p < q$ if the path p differs from p_{m+1} by fewer tiles than q . The minimal path under this order is of course p_{m+1} , from which we obtain the image under ϕ of d_{m+1} . Now suppose all basis elements obtained from paths that differ from p_{m+1} by at most r tiles are in the image of ϕ , and choose a path p that differs by $r+1$. Then p must be obtained from p_{m+1} by adding a (minimal) ordered sequence of $r+1$ tiles at positions $i_1, i_2, \dots, i_{r+1}$. Consider now the diagram $d = e_{i_{r+1}} \dots e_{i_2} e_{i_1} d_{m+1}$, which is mapped under ϕ to a path

$$\begin{aligned} \phi(d) &= e_{i_{r+1}} \dots e_{i_2} e_{i_1} \phi(d_{m+1}) \\ &= e_{i_{r+1}} \dots e_{i_2} e_{i_1} v_{p_{m+1}}. \end{aligned}$$

By Theorem 4.5, we see that this must equal

$$v_p + \sum_{q < p} \alpha_q v_q$$

where p is the original path, each path q differs from p_{m+1} by r or fewer tiles, and α_q is a constant. By induction each q is the image of an element x_q of $W_{+,+}^{(n,m)}$, hence v_p is the image of $d - \sum_{q < p} \alpha_q x_q$.

Finally, note that the dimensions of the modules are equal [5, Theorem 6.4], therefore the homomorphism ϕ must in fact be an isomorphism.

For $\varepsilon_1 = -1, \varepsilon_2 = 1$ we repeat the above argument by mapping the path of final height $-m - 1$ which is closest to the fundamental path to the diagram d_{-m-1} in (22).

$$d_{-m-1} = \begin{array}{c} \begin{array}{|c|c|c|c|c|} \hline \bullet & \bullet & \dots & \bullet & \dots \\ \hline \bullet & \bullet & \dots & \bullet & \dots \\ \hline \end{array} \\ \underbrace{\hspace{1.5cm}}_m \end{array} \quad (22)$$

If now we choose $\varepsilon_1 = 1, \varepsilon_2 = -1$, then we map the path of final height $m + 1$ which is closest to the fundamental path to the diagram d'_{m+1} in (23).

$$d'_{m+1} = \begin{array}{c} \begin{array}{|c|c|c|c|c|} \hline \bullet & \bullet & \dots & \bullet & \dots \\ \hline \bullet & \bullet & \dots & \bullet & \dots \\ \hline \end{array} \\ \underbrace{\hspace{1.5cm}}_m \end{array} \quad (23)$$

For $\varepsilon_1 = -1, \varepsilon_2 = -1$ we repeat the above argument by mapping the path of final height $-m - 1$ which is closest to the fundamental path to the diagram d'_{-m-1} in (24).

$$d'_{-m-1} = \begin{array}{c} \begin{array}{|c|c|c|c|c|} \hline \bullet & \bullet & \dots & \bullet & \dots \\ \hline \bullet & \bullet & \dots & \bullet & \dots \\ \hline \end{array} \\ \underbrace{\hspace{1.5cm}}_{m-1} \end{array} \quad (24)$$

□

Corollary 6.9. *Fix δ and hence b_n^x . Whenever $\alpha_{\varepsilon_1, \varepsilon_2}^{(n, m)} \in k$, the central element Z_n acts by $\alpha_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ on $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$.*

Proof. Use [5, Theorem 6.4] and the isomorphism from Theorem 6.8 for the generic case. The other cases follow by analytic continuity — Z_n acts by some scalar, so the limit of generic actions approaching δ exists and is this scalar. □


An important and immediate consequence is the following corollary.

Corollary 6.10. *Fix δ and hence b_n^x . A necessary condition for two cell modules $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ and $W_{\eta_1, \eta_2}^{(n, t)}$ to be in the same block is that $\alpha_{\varepsilon_1, \varepsilon_2}^{(n, m)} = \alpha_{\eta_1, \eta_2}^{(n, t)}$.*

7. GRAM DETERMINANTS FOR CELL MODULES

We recall from [7, §8.2] that each cell module $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ has a basis $B_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ of half diagrams with arcs on the northern edge (hereafter referred to as the standard diagram basis).

Example 7.1. The cell module $W_{-, -}^{(5, 2)}$ has the following basis of half diagrams:



Consider $u, v \in B_{\varepsilon_1, \varepsilon_2}^{(n, m)}$. We define a scalar $\langle u, v \rangle$ as follows. We first flip v vertically and identify the southern nodes of this diagram with the respective northern nodes of u . After applying the straightening rules for b_n^x , we obtain a diagram with a number of (possibly decorated) strings. The

value of $\langle u, v \rangle$ is the coefficient of this diagram if the strings match the number and decorations needed for the cell module, and is zero otherwise. For instance

$$\begin{array}{c} \text{Diagram 1} \end{array} = \delta_L^2 \delta_R \begin{array}{c} \text{Diagram 2} \end{array} \quad \text{and} \quad \begin{array}{c} \text{Diagram 3} \end{array} = \delta_L \begin{array}{c} \text{Diagram 4} \end{array},$$

giving $\langle u, v \rangle = \delta_L^2 \delta_R$ and $= 0$ respectively. (This is $\langle u, v \rangle$ as defined in [12, §6] with the “semime-ander” convention.)

Proposition 7.2. *Let $\sigma : b_n^x \rightarrow b_n^x$ be the involution defined by flipping diagrams vertically. Then the inner product defined by $\langle -, - \rangle$, together with σ , defines a contravariant bilinear form on $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$. That is, for $d \in b_n^x$, we have*

$$\langle du, v \rangle = \langle u, \sigma(d)v \rangle.$$

Proof. This follows from the definition of $\langle u, v \rangle$ and the action of the algebra on half diagrams. $\square$

While this form can be used over the integral ring $\mathbb{Z}[q^\pm, Q_1^\pm, Q_2^\pm]$, we will need to specialise in order to use results from the parameterisation of De Gier–Nichols.

We define the Gram matrix $G_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ to be the matrix of entries $(\langle u, v \rangle)_{u, v}$, where u, v runs over the basis $B_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ of $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$. We also define the Gram determinant

$$\Gamma_{\varepsilon_1, \varepsilon_2}^{(n, m)} = \det G_{\varepsilon_1, \varepsilon_2}^{(n, m)}. \quad (25)$$

When we base change to a field, the rank of $G_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ is also the rank of a corresponding map from the module to its contravariant dual. The module is thus simple if and only if the matrix is non-singular.

Example 7.1 contd. With the ordering of basis elements as in 7.1, the Gram matrix is therefore

$$G_{-, -}^{(5, 2)} = \begin{pmatrix} \delta_L^2 \delta_R \kappa_L & \delta_L \delta_R \kappa_L & \delta_L^2 \delta_R & 0 & 0 & 0 \\ \delta_L \delta_R \kappa_L & \delta_L \delta_R \delta & \delta_L \delta_R & 0 & 0 & 0 \\ \delta_L^2 \delta_R & \delta_L \delta_R & \delta_L \delta_R \delta & \delta_L \delta_R & 0 & 0 \\ 0 & 0 & \delta_L \delta_R & \delta_L \delta_R \delta & \delta_L \delta_R & \delta_L \delta_R^2 \\ 0 & 0 & 0 & \delta_L \delta_R & \delta_L \delta_R \delta & \delta_L \delta_R \kappa_R \\ 0 & 0 & 0 & \delta_L \delta_R^2 & \delta_L \delta_R \kappa_R & \delta_L \delta_R^2 \kappa_R \end{pmatrix}.$$

We wish to calculate the determinant of this matrix. By Laplace expansion, we obtain

$$(\delta_L \delta_R)^{-6} \Gamma_{-, -}^{(5, 2)} = \delta_L \kappa_L \begin{vmatrix} \delta & 1 & 0 & 0 & 0 \\ 1 & \delta & 1 & 0 & 0 \\ 0 & 1 & \delta & 1 & \delta_R \\ 0 & 0 & 1 & \delta & \kappa_R \\ 0 & 0 & \delta_R & \kappa_R & \delta_R \kappa_R \end{vmatrix} - \kappa_L^2 \begin{vmatrix} \delta & 1 & 0 & 0 \\ 1 & \delta & 1 & \delta_R \\ 0 & 1 & \delta & \kappa_R \\ 0 & \delta_R & \kappa_R & \delta_R \kappa_R \end{vmatrix} + (2\delta_L \kappa_L - \delta_L^2 \delta) \begin{vmatrix} \delta & 1 & \delta_R \\ 1 & \delta & \kappa_R \\ \delta_R & \kappa_R & \delta_R \kappa_R \end{vmatrix}.$$

Laplace expanding the first of these determinants results in the following:

$$\delta_R \kappa_R \begin{vmatrix} \delta & 1 & 0 & 0 \\ 1 & \delta & 1 & 0 \\ 0 & 1 & \delta & 1 \\ 0 & 0 & 1 & \delta \end{vmatrix} - \kappa_R^2 \begin{vmatrix} \delta & 1 & 0 \\ 1 & \delta & 1 \\ 0 & 1 & \delta \end{vmatrix} + (2\delta_R \kappa_R - \delta \delta_R^2) \begin{vmatrix} \delta & 1 \\ 1 & \delta \end{vmatrix}. \quad (26)$$

Since $\delta = [2]$, we can use the identities for quantum integers to show that each of these determinants is equal to $[n+1]$, where n is the size of the matrix. Using our parameterisation for $\delta_L, \delta_R, \kappa_L, \kappa_R$, we then see that (26) is equal to

$$\begin{aligned}
& [w_2 + 1]^{-2} ([w_2][w_2 + 1][5] - [w_2 + 1]^2[4] + (2[w_2][w_2 + 1] - [2][w_2]^2)[3]) \\
&= [w_2 + 1]^{-2} ([w_2][w_2 + 1][5] - [w_2 + 1]^2[4] + [w_2]([w_2 + 1] - [w_2 - 1])[3]) \\
&= [w_2 + 1]^{-2} ([w_2]([w_2 + 1][5] - [w_2 - 1][3]) - [w_2 + 1]([w_2 + 1][4] - [w_2][3])) \\
&= [w_2 + 1]^{-2} ([w_2][2][w_2 + 4] - [w_2 + 1][w_2 + 4]) \\
&= [w_2 + 1]^{-2} [w_2 + 4][w_2 - 1].
\end{aligned}$$

Expanding the other matrices in the same way, we see that

$$\begin{aligned}
& \left(\frac{[w_1][w_2]}{[w_1 + 1][w_2 + 1]} \right)^{-6} [w_1 + 1]^2 [w_2 + 1]^2 G_{-, -}^{(5, 2)} \\
&= [w_1][w_1 + 1][w_2 + 4][w_2 - 1] - [w_1 + 1]^2 [w_2 + 3][w_2 - 1] \\
&\quad + (2[w_1][w_1 + 1] - [w_1]^2[2]) [w_2 + 2][w_2 - 1] \\
&= [w_2 - 1] \left([w_1][w_1 + 1][w_2 + 4] - [w_1 + 1]^2 [w_2 + 3] \right. \\
&\quad \left. + [w_1]([w_1 + 1] - [w_1 - 1])[w_2 + 2] \right).
\end{aligned}$$

Note that the last expression in the brackets takes the same form as the second line in our evaluation of (26) above. Hence we finally arrive at

$$\Gamma_{-, -}^{(5, 2)} = \frac{[w_1]^6 [w_2]^6}{[w_1 + 1]^8 [w_2 + 1]^8} [w_1 - 1][w_2 - 1][w_1 + w_2 + 3].$$

As demonstrated by this example, calculating $\Gamma_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ is non-trivial. However we can apply results of [5] to calculate it with respect to the path basis, which we will see is easier.

Proposition 7.3 ([5, Proposition 5.13]). *In the path basis, $\hat{G}_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ is diagonal.*

Definition 7.4 ([5, Definition 5.14]). We define the functions $f(h)$ and $g(h)$ to be:

$$\begin{aligned}
f(h) &= r(w_1 - h)r(-w_1 + h) \\
g(h) &= k(w_1 - h)k(-w_1 + h)
\end{aligned}$$

where $r(u)$ and $k(u)$ are as in (6).


Proposition 7.5 ([5, Proposition 5.15]). *The eigenvalue λ_p of the Gram matrix for each path p is given by the following recursive procedure. Let p_0 be the fundamental path, and let p' be a path obtained from another path p by the addition of a tile (or half tile) at point i . The following hold:*

- $\lambda_{p_0} = 1$.
- If p' and p differ by a full tile we have $\lambda_{p'} = f(h_{i-1})\lambda_p$.

- If p' and p differ by a half tile we have $\lambda_{p'} = g(h_{n-1})\lambda_p$.

Thus to find $\Gamma_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ we take the product of the eigenvalues corresponding to the paths that form a basis of that cell module (having chosen θ appropriately). To illustrate, we return to Example 7.1 above, and recalculate the Gram matrix with respect to the path basis.

Example 7.6. The basis of $W_{-, -}^{(5, 2)}$ here consists of all paths of a final height -3 or lower, and our value of θ is $-2 - w_1 - w_2$. These paths are given below, along with the tiles that are needed to construct them.



The eigenvalues of the Gram matrix for these paths are

$$1, f(0), f(-1), f(-1)f(-2), f(-1)f(-2)f(-3), f(-1)f(-2)f(-3)g(-4)$$

respectively. The Gram determinant is the product of these, which we evaluate to be

$$\Gamma_{-, -}^{(5, 2)} = \frac{[w_1]^2}{[w_1 + 1]^4 [w_2 + 1]^2} [w_1 - 1] [w_2 - 1] [w_1 + w_2 + 3].$$

Note that this is the same result as Example 7.1, up to rescaling by a power of the parameters.

Note this is easier than the calculation in Example 7.1. However in general $\Gamma_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ may still be difficult to calculate, due to the large number of paths and tiles as the cell modules increase in size. We next appeal to results about changing bases and the effect on the Gram determinant.

Theorem 7.7. *With respect to the path basis above, the Gram determinant of $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ is*

$$\begin{aligned} \Gamma_{\varepsilon_1, \varepsilon_2}^{(n, m)} &= (\delta_L^{\frac{1}{2}(1-\varepsilon_1)} \delta_R^{\frac{1}{2}(1-\varepsilon_2)})^{\dim W_{\varepsilon_1, \varepsilon_2}^{(n, m)}} \prod_{k=0}^{\frac{1}{2}(n-m-3)} \left(\left[\frac{1}{2}(n-m-2k-1) \right] \right. \\ &\quad \times [\varepsilon_1 w_1 - \frac{1}{2}(-n+m+2k+1)] [\varepsilon_2 w_2 - \frac{1}{2}(-n+m+2k+1)] \\ &\quad \left. \times [\varepsilon_1 w_1 + \varepsilon_2 w_2 - \frac{1}{2}(n+m-2k-1)] [w_1 + 1]^{-2} [w_2 + 1]^{-2} \right)^{\dim W_{\varepsilon_1, \varepsilon_2}^{(n, n-1-2k)}}. \end{aligned}$$

Proof. From the definitions of $f(h)$ and $g(h)$, we see that $\Gamma_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ is a product of box numbers of the form $[a_1 w_1 + a_2 w_2 - b]^c$, where $a_i \in \{-1, 0, 1\}$ and $b, c \in \mathbb{Z}$. Note that all such terms with either $a_1 = 0$ or $a_1, a_2 \neq 0$ arise from the contributions of some $g(h)$ at the right boundary. Moreover, apart from $[w_2 + 1]$ they all appear to a positive power. Therefore when calculating the product over all permitted paths, there can be no cancellation of these terms. To determine the power of $[w_2 + 1]$, we multiply the above product by $[w_2 + 1]^{-2}$ for every factor $g(h)$. The power to which

$g(h)$ appears in the product is the number of paths of final height h' for $|h'| \geq |h|$, which in turn is the dimension of the cell module defined by paths of such height. Therefore we see that

$$\mu \prod_{k=0}^{\frac{1}{2}(n-m-3)} \left(\left[\frac{1}{2}(n-m-2k-1) \right] [\varepsilon_2 w_2 - \frac{1}{2}(-n+m+2k+1)] \right. \\ \left. \times [\varepsilon_1 w_1 + \varepsilon_2 w_2 - \frac{1}{2}(n+m-2k-1)] [w_2 + 1]^{-2} \right)^{\dim W_{\varepsilon_1, \varepsilon_2}^{(n, n-1-2k)}}$$

is a factor of $\Gamma_{\varepsilon_1, \varepsilon_2}^{(n, m)}$, where μ is a product of box numbers of the form $[w_1 - a]$ for $a \in \mathbb{Z}$.

In order to determine the other factors, we will change basis and recalculate the Gram determinant. First, note from the proof of Theorem 6.8 that the change of basis matrix between the standard and path bases is upper triangular, with diagonal entries equal to powers of the parameters for the symplectic blob algebra. Note also that these diagonal entries do not contain the parameter δ . Indeed, the relations that could result in a factor of δ must be the standard Temperley-Lieb relations, i.e.

$$e_i e_{i \pm 1} e_i = e_i \text{ and } e_i^2 = \delta e_i,$$

but these cannot appear in the leading term of the path basis as we can never add tiles in position i , followed by $i \pm 1$, then in i again, nor can we add tiles in position i twice in a row. We also cannot obtain a δ by adding to the initial diagram d_{m+1} (or $d_{-m-1}, d'_{m+1}, d'_{-m-1}$).

From the standard diagram basis, we change to an alternative path basis, which we obtain by replacing e_i by e_{n-i} , w_1 by w_2 and ε_1 by ε_2 in the above. In other words, we are working with the path basis defined by the right blob as opposed to the left. For the same reasons as in Theorem 6.8, the change of basis matrix is again upper triangular. Therefore the change of basis matrix between the first and second path bases is upper triangular, and has determinant equal to a product of powers of the parameters (except δ , as before). Moreover, by considering the contribution at the half tile boundary in the second path basis, we see that

$$\mu' \prod_{k=0}^{\frac{1}{2}(n-m-3)} \left(\left[\frac{1}{2}(n-m-2k-1) \right] [\varepsilon_1 w_1 - \frac{1}{2}(-n+m+2k+1)] \right. \\ \left. \times [\varepsilon_1 w_1 + \varepsilon_2 w_2 - \frac{1}{2}(n+m-2k-1)] [w_1 + 1]^{-2} \right)^{\dim W_{\varepsilon_1, \varepsilon_2}^{(n, n-1-2k)}}$$

is a factor of $\Gamma_{\varepsilon_1, \varepsilon_2}^{(n, m)}$, where μ' is a product of powers of the parameters and box numbers of the form $[w_2 - a]$ for $a \in \mathbb{Z}$. When we combine these two results we have

$$\Gamma_{\varepsilon_1, \varepsilon_2}^{(n, m)} = \lambda \prod_{k=0}^{\frac{1}{2}(n-m-3)} \left(\left[\frac{1}{2}(n-m-2k-1) \right] [\varepsilon_1 w_1 - \frac{1}{2}(-n+m+2k+1)] \right. \\ \left. \times [\varepsilon_2 w_2 - \frac{1}{2}(-n+m+2k+1)] [\varepsilon_1 w_1 + \varepsilon_2 w_2 - \frac{1}{2}(n+m-2k-1)] [w_1 + 1]^{-2} [w_2 + 1]^{-2} \right)^{\dim W_{\varepsilon_1, \varepsilon_2}^{(n, n-1-2k)}},$$

where λ is a product of powers of the parameters δ_L, δ_R (since our parameterisation has $\kappa_L = \kappa_R = 1$). To determine λ , we return to the Gram matrix of the standard diagram basis and determine the highest powers of δ_L and δ_R which divide the determinant. Since propagating lines cannot cross, any non-zero entry in the Gram matrix must have a factor of δ_L (resp. δ_R) if there is a left (resp. right) blob on propagating lines. Therefore we can extract a factor of $(\delta_L^{\frac{1}{2}(1-\varepsilon_1)} \delta_R^{\frac{1}{2}(1-\varepsilon_2)})^{\dim W_{\varepsilon_1, \varepsilon_2}^{(n, m)}}$ from the matrix. In fact, this is the largest power of δ_L and δ_R we can extract from any row. We deal with factors that may arise from further left blobs, those arising from the right follow by symmetry. Suppose a diagram has a horizontal arc with a left blob. Then this must be the outermost arc of a left-exposed nest of arcs. We construct a diagram which, when taking the inner product with the first, does not add any factors of δ_L with this nested set of arcs. Indeed, we simply place undecorated arcs in the leftmost side of the diagram so that the blobbed arc forms a closed loop after taking the inner product. This results in a factor of $\kappa_L = 1$ appearing, and no δ_L .

Finally, the range of values over which we take the product ensure that neither $[w_1]$ nor $[w_2]$ can appear. Therefore λ must be the greatest factor of δ_L and δ_R , and the result follows. $\square$

This final example returns to the cell module $W_{-, -}^{(5, 2)}$.

Example 7.8. For $n = 5, m = 2$, we have $\frac{1}{2}(n - m - 3) = 0$. Therefore by Theorem 7.7, we have

$$\begin{aligned} \Gamma_{-, -}^{(5, 2)} &= ([w_1][w_1 + 1]^{-1}[w_2][w_2 + 1]^{-1})^6 [w_1 + 1]^{-2} [w_2 + 1]^{-2} [1][-w_1 + 1][-w_2 + 1][-w_1 - w_2 - 3] \\ &= [w_1]^6 [w_1 + 1]^{-8} [w_2]^6 [w_2 + 1]^{-8} [w_1 - 1][w_2 - 1][w_1 + w_2 + 3]. \end{aligned}$$

We can compare this with Example 7.1 to see that we indeed have the Gram determinant.

8. HOMOLOGICAL TOOLS FOR DECOMPOSITION MATRICES AND BLOCKS OF b'_n

In this section, we will use the constants $\alpha_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ and homomorphisms from [9] to determine the block structure of b_n^x for $\delta \in \mathbb{C}^6$. Fixing δ is done by choosing values for q, w_1 and w_2 . We will here restrict those values such that none of $[w_1], [w_2], [w_1 + 1]$ or $[w_2 + 1]$ are zero. We now change our parameterisation to that of GMP2 in §1 above, in order to use the results of [9]. This is achieved by rescaling generators in the following way:

$$\begin{aligned} e_0 &\mapsto -[w_1 + 1]e_0, \\ e_i &\mapsto -e_i \text{ for } 1 \leq i \leq n - 1 \\ e_n &\mapsto -[w_2 + 1]e_n. \end{aligned}$$

8.1. Globalisation functors. We will also use the globalisation functors to work in a “large n limit” symplectic blob algebra where both parameters are positive. Having determined blocks in this limit, we will then localise back to the original algebra with original parameter values. The following proposition taken from [9, §3] justifies this.

Proposition 8.1 ([9, §3]). *There exist right exact globalisation functors*

$$G : b_n^x - \text{mod} \longrightarrow b_{n+1}^x - \text{mod}, \quad G' : b_n^x - \text{mod} \longrightarrow b_{n+1}^x - \text{mod}$$

with the following properties:

- (1) *There is a parameter change from b_n^x to b_{n+1}^x under G which sends $w_1 \mapsto -w_1 - 1$;*
- (2) *There is a parameter change under G' which sends $w_2 \mapsto -w_2 - 1$;*
- (3) $GW_{\varepsilon_1, \varepsilon_2}^{(n, m)} = W_{-\varepsilon_1, \varepsilon_2}^{(n+1, m+\varepsilon_1)}$;
- (4) $G'W_{\varepsilon_1, \varepsilon_2}^{(n, m)} = W_{\varepsilon_1, -\varepsilon_2}^{(n+1, m+\varepsilon_2)}$.

There are also exact localisation functors

$$F : b_n^x - \text{mod} \longrightarrow b_{n-1}^x - \text{mod}$$

$$F' : b_n^x - \text{mod} \longrightarrow b_{n-1}^x - \text{mod}$$

such that $F \circ G = \text{id}$ and $F' \circ G' = \text{id}$, and also

$$(1) \quad FW_{\varepsilon_1, \varepsilon_2}^{(n, m)} = \begin{cases} W_{-\varepsilon_1, \varepsilon_2}^{(n-1, m+\varepsilon_1)} & \text{if } m \neq n-1 \text{ or } \varepsilon_1 = -1 \\ 0 & \text{if } m = n-1 \text{ and } \varepsilon_1 = 1; \end{cases}$$

$$(2) \quad F'W_{\varepsilon_1, \varepsilon_2}^{(n, m)} = \begin{cases} W_{\varepsilon_1, -\varepsilon_2}^{(n-1, m+\varepsilon_2)} & \text{if } m \neq n-1 \text{ or } \varepsilon_2 = -1 \\ 0 & \text{if } m = n-1 \text{ and } \varepsilon_2 = 1. \end{cases}$$

Note that the localisation functor can annihilate modules, and therefore it is possible for a block to “break up” when localising. We will address this on a case by case basis when determining the blocks below. Also, since we will always localise back after globalising, we need only consider in the arguments below cell modules $W_{\varepsilon_1, \varepsilon_2}^{(N, m)}$ with $m \ll N$.

8.2. On standard module homomorphisms. We now recall the homomorphisms from [9] and reformulate them into the notation consistent with this paper.

Theorem 8.2 ([9, Theorem 4.3.5]). *Let q be a primitive 2ℓ -th root of unity and $w_1, w_2 \notin \mathbb{Z}$. Suppose that $m - 2\ell \geq 0$ (with equality if and only if $\varepsilon_1 = \varepsilon_2 = 1$). Then there exists a non-zero homomorphism*

$$\psi : W_{\varepsilon_1, \varepsilon_2}^{(n, m)} \longrightarrow W_{\varepsilon_1, \varepsilon_2}^{(n, m-2\ell)}.$$

Theorem 8.3 ([9, Theorem 4.1.3]). *Let q be a primitive 2ℓ -th root of unity and $w_1 \in \mathbb{Z}$. Suppose for $r \in \mathbb{Z}$ that $m > m - 2(\varepsilon_1 w_1 + r\ell) > 0$. Then there exists a non-zero homomorphism*

$$\psi : W_{\varepsilon_1, \varepsilon_2}^{(n, m)} \longrightarrow W_{-\varepsilon_1, \varepsilon_2}^{(n, m-2(\varepsilon_1 w_1 + r\ell))}.$$

If q is not a root of unity, then set $\ell = 0$ in the above.

Theorem 8.4 ([9, Theorems 4.1.6 and 4.1.7]). *Let q be a primitive 2ℓ -th root of unity and $w_2 \in \mathbb{Z}$. Suppose for $r \in \mathbb{Z}$ that $m > m - 2(\varepsilon_2 w_2 + r\ell) > 0$. Then there exists a non-zero homomorphism*

$$\psi : W_{\varepsilon_1, \varepsilon_2}^{(n, m)} \longrightarrow W_{\varepsilon_1, -\varepsilon_2}^{(n, m - 2(\varepsilon_2 w_2 + r\ell))}.$$

If q is not a root of unity then set $\ell = 0$ in the above.

Theorem 8.5 ([9, Theorems 4.2.11 and 4.2.12]). *Let q be a primitive 2ℓ -th root of unity and $\varepsilon_1 w_1 + \varepsilon_2 w_2 \in \mathbb{Z}$. Suppose for $r \in \mathbb{Z}$ that $m > 2(\varepsilon_1 w_1 + \varepsilon_2 w_2 + r\ell) - m \geq 0$ (with equality only if $\varepsilon_1 = \varepsilon_2 = 1$). Then there exists a non-zero homomorphism*

$$\psi : W_{\varepsilon_1, \varepsilon_2}^{(n, m)} \longrightarrow W_{\varepsilon_1, \varepsilon_2}^{(n, 2(\varepsilon_1 w_1 + \varepsilon_2 w_2 + r\ell) - m)}.$$

If q is not a root of unity then set $\ell = 0$ in the above.

8.3. Block master equations. By Proposition 6.4 a necessary condition for any two cell modules to be in the same block is that Z_n acts by the same constant on both modules. Notice that

$$\begin{aligned} \alpha_{\varepsilon_1, \varepsilon_2}^{(n, m)} &= [n] \frac{[2(-m + \varepsilon_1 w_1 + \varepsilon_2 w_2)]}{[-m + \varepsilon_1 w_1 + \varepsilon_2 w_2]} \\ &= [n] (q^{-m + \varepsilon_1 w_1 + \varepsilon_2 w_2} + q^{m - \varepsilon_1 w_1 - \varepsilon_2 w_2}) \end{aligned}$$

Therefore if $\alpha_{\varepsilon_1, \varepsilon_2}^{(n, m)} = \alpha_{\eta_1, \eta_2}^{(n, t)}$ and $[n] \neq 0$, then

$$q^{-m + \varepsilon_1 w_1 + \varepsilon_2 w_2} + q^{m - \varepsilon_1 w_1 - \varepsilon_2 w_2} = q^{-t + \eta_1 w_1 + \eta_2 w_2} + q^{t - \eta_1 w_1 - \eta_2 w_2}.$$

Thus we have $q^{-m + \varepsilon_1 w_1 + \varepsilon_2 w_2} = q^{\pm(-t + \eta_1 w_1 + \eta_2 w_2)}$. This can only be satisfied if either

$$-(m - t) + (\varepsilon_1 - \eta_1)w_1 + (\varepsilon_2 - \eta_2)w_2 \equiv 0 \pmod{2\ell}, \text{ or}$$

$$-(m + t) + (\varepsilon_1 + \eta_1)w_1 + (\varepsilon_2 + \eta_2)w_2 \equiv 0 \pmod{2\ell}.$$

In the first case, the allowed values of η_1, η_2 lead to the following possibilities:

$$\varepsilon_1 \neq \eta_1, \varepsilon_2 \neq \eta_2 \implies m - t \equiv 2\varepsilon_1 w_1 + 2\varepsilon_2 w_2 \pmod{2\ell} \quad (27)$$

$$\varepsilon_1 \neq \eta_1, \varepsilon_2 = \eta_2 \implies m - t \equiv 2\varepsilon_1 w_1 \pmod{2\ell} \quad (28)$$

$$\varepsilon_1 = \eta_1, \varepsilon_2 \neq \eta_2 \implies m - t \equiv 2\varepsilon_2 w_2 \pmod{2\ell} \quad (29)$$

$$\varepsilon_1 = \eta_1, \varepsilon_2 = \eta_2 \implies m - t \equiv 0 \pmod{2\ell}, \quad (30)$$

and in the second case we have:

$$\varepsilon_1 = \eta_1, \varepsilon_2 = \eta_2 \implies m + t \equiv 2\varepsilon_1 w_1 + 2\varepsilon_2 w_2 \pmod{2\ell} \quad (31)$$

$$\varepsilon_1 = \eta_1, \varepsilon_2 \neq \eta_2 \implies m + t \equiv 2\varepsilon_1 w_1 \pmod{2\ell} \quad (32)$$

$$\varepsilon_1 \neq \eta_1, \varepsilon_2 = \eta_2 \implies m + t \equiv 2\varepsilon_2 w_2 \pmod{2\ell} \quad (33)$$

$$\varepsilon_1 \neq \eta_1, \varepsilon_2 \neq \eta_2 \implies m + t \equiv 0 \pmod{2\ell}. \quad (34)$$




FIGURE 6. Graphical depiction of the cell modules of b'_8 with $w_1 = \frac{1}{2}$ and $w_2 = \frac{3}{4}$.

If q is not a root of unity, then all of the congruences modulo 2ℓ in the above become equalities.

If $[n] = 0$, then we can still use equations (27)–(34) by first globalising to b_N^x where $[N] \neq 0$, determining the blocks there, and then localising again.

9. DECOMPOSITION MATRICES AND BLOCKS OF b'_n

In the following subsections we will consider separately various cases relating to whether or not certain linear combinations of w_1 and w_2 are integers.

To visualise solutions to the master equations (27–34), we will plot points in the plane corresponding to cell modules, in such a way that solutions are manifested geometrically. (Remark: this indicates the potential for a geometric linkage principle, cf. [13], to describe the representation theory of the algebra.) The cell module $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ is given ‘weight’ coordinates

$$W_{\varepsilon_1, \varepsilon_2}^{(n, m)} \mapsto (\varepsilon_1(m - \varepsilon_1 w_1 - \varepsilon_2 w_2), \varepsilon_2(m - \varepsilon_1 w_1 - \varepsilon_2 w_2)),$$

— see e.g. Figure 6, Figure 9. In this geometry (in the q not a root of unity case) two cell modules have the same Z_n -eigenvalue if and only if one can be reached from the other by successive reflections in the coordinate axes. As a guide to the eye, the cell module $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ is plotted on the ‘arm’ labelled by $\varepsilon_1, \varepsilon_2$.

9.1. Cases with none of $w_1, w_2, w_1 + w_2, w_1 - w_2$ integral. Suppose first that q is not a root of unity. Since m and t are positive integers, it is only possible for at most one of (27) and (31) to be satisfied (similarly for (28) and (32); (29) and (33); and (30) and (34)). The case (34) is impossible as both m and t are positive integers, and at most one can be zero. The case (30) is trivial, as the two modules are equal here. Also since ε_1 and ε_2 take values ± 1 , we can have non-trivial coincidences of the eigenvalues of Z_n if and only if $\{w_1, w_2, w_1 + w_2, w_1 - w_2\} \cap \mathbb{Z} \neq \emptyset$. This leads to the first main theorem of this paper:

Theorem 9.1. *Suppose q is not a root of unity and $\{w_1, w_2, w_1 + w_2, w_1 - w_2\} \cap \mathbb{Z} = \emptyset$. Then the algebra b'_n is semisimple. If in addition, $\theta \neq \pm(-m \pm w_1 \pm w_2)$ for any $m \in \mathbb{Z}$ then symplectic blob algebra b_n^x is semisimple.*

Proof. To prove the first statement, it suffices to show that the eigenvalues of Z_n are all distinct. Indeed, since none of $w_1, w_2, w_1 + w_2$ or $w_1 - w_2$ are integral the only possible solution to equations (27)–(34) is the trivial one in (30). Therefore each cell module is alone in its block and the algebra is semisimple.

To prove the second, the only additional information needed is that $W^{(n)}(b)$ is simple. This is guaranteed as for our chosen value of θ , the Gram determinant of $W^{(n)}(b)$ is non-zero by [5, Theorem 5.17]. $\square$

If now q is a 2ℓ -th root of unity, then we must consider equations (30) and (34). Note that the left- and right-blob algebras are semisimple, so if $\eta_1 \neq \varepsilon_1$ and $\eta_2 \neq \varepsilon_2$ then by restricting to either algebra and considering the standard contents of Table 4 we see that there can be no homomorphisms between any modules satisfying (34). It therefore remains to consider (30). Suppose without loss of generality that $t < m$. Then $0 \leq m - 2\ell$ (with equality only if $\varepsilon_1 = \varepsilon_2 = 1$), so by Theorem 8.2, we have a non-zero homomorphism

$$W_{\varepsilon_1, \varepsilon_2}^{(n, m)} \longrightarrow W_{\varepsilon_1, \varepsilon_2}^{(n, m-2\ell)} \quad (35)$$

Thus $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ and $W_{\varepsilon_1, \varepsilon_2}^{(n, m-2\ell)}$ are in the same block. Similarly we have $W_{\varepsilon_1, \varepsilon_2}^{(n, m-2\ell)}$ and $W_{\varepsilon_1, \varepsilon_2}^{(n, m-4\ell)}$ in the same block, and so on. By transitivity, we therefore see that $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ and $W_{\varepsilon_1, \varepsilon_2}^{(n, t)}$ are in the same block.

Theorem 9.2. *Suppose $q^{2\ell} = 1$ and $\{w_1, w_2, w_1 + w_2, w_1 - w_2\} \cap \mathbb{Z} = \emptyset$. Then two cell modules $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ and $W_{\eta_1, \eta_2}^{(n, t)}$ are in the same block if and only if $\varepsilon_1 = \eta_1$, $\varepsilon_2 = \eta_2$ and $m \equiv t \pmod{2\ell}$.*

The combinatorial-geometric expression of linkage in this case is as in Figure 7. See Figure 8 for the truncation to $n = 8$.

9.2. Either w_1 or w_2 integral. We will determine the blocks when precisely one of w_1 and w_2 is integral. We begin with the case $w_1 \in \mathbb{Z}, w_2 \notin \mathbb{Z}$, and first assume that q is not a root of unity. Now the only equations from (27)–(34) with non-trivial solutions are (28) and (32), and by fixing m, ε_1 and ε_2 we see that blocks have size at most two. Note that if $w_2 \in \frac{1}{2}\mathbb{Z}$ then we still do not obtain extra solutions since $m \pm t$ is always even.

Consider first the case (32), where we have $\varepsilon_1 = \eta_1$ and $\varepsilon_2 \neq \eta_2$. We will show that although these two modules have the same eigenvalue, they are not in the same block. As right-blob modules, they have a filtration as in Table 4. However since the parameter w_2 is not integral the right-blob algebra is semisimple, and thus it is not possible to have a non-zero homomorphism between the




FIGURE 7. Cell modules of b'_{14} with $w_1 = \frac{1}{2}$ and $w_2 = \frac{3}{4}$. Dashed lines indicate modules in the same block when $\ell = 3$.




FIGURE 8. Graphical depiction of the cell modules of b'_8 with $w_1 = \frac{1}{2}$ and $w_2 = \frac{3}{4}$. The dashed lines indicate modules in the same block when $\ell = 3$.

modules. Since the block has size at most two, we deduce that these modules are not in the same block.

Now consider (28). Here, we have $\varepsilon_1 \neq \eta_1$ and $\varepsilon_2 = \eta_2$. By swapping labels if necessary we may assume that $m > t$ (equality is not possible due to [9, Proposition 3.4.1]). Since both m and t are non-negative integers we must have $\varepsilon_1 = \text{sgn}(w_1)$, thus $m > t = m - 2\varepsilon_1 w_1 > 0$. Therefore the conditions of Theorem 8.3 are satisfied and we have a homomorphism $W_{\varepsilon_1, \varepsilon_2}^{(n, m)} \longrightarrow W_{-\varepsilon_1, \varepsilon_2}^{(n, t)}$.




FIGURE 9. Graphical depiction of the cell modules of (i) b'_8 with $w_1 = 1$ and $w_2 = \frac{3}{4}$; and (ii) b'_9 with $w_1 = -\frac{1}{4}$ and $w_2 = 1$.

We will now consider the case when q is a 2ℓ -th root of unity and $w_1 \in \mathbb{Z}, w_2 \notin \mathbb{Z}$. In this case, the only equations from (27)–(34) with solutions are (28), (30), (32) and (34). We still do not obtain extra solutions if $w_2 \in \frac{1}{2}\mathbb{Z}$ by parity considerations in the same way as above.

Begin by fixing the cell module with labels m, ε_1 and ε_2 . By restricting to the right-blob algebra as before, any other cell module $W_{\eta_1, \eta_2}^{(n, t)}$ in this block has $\eta_2 = \varepsilon_2$. We can therefore rule out equations (32) and (34). In the case of equation (28) we again see that the conditions of Theorem 8.3 are satisfied (this time with $\ell \neq 0$), and so these cell modules are in the same block. So it remains to consider the case of (30). We will begin by showing that $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ and $W_{\varepsilon_1, \varepsilon_2}^{(n, m+2\ell)}$ are in the same block, and the general result will follow. Indeed, we choose $r \in \mathbb{Z}$ such that $0 < \varepsilon_1 w_1 + r\ell < \ell$, then by Theorem 8.3 we have non-zero homomorphisms $W_{\varepsilon_1, \varepsilon_2}^{(n, m+2\ell)} \rightarrow W_{-\varepsilon_1, \varepsilon_2}^{(n, m+2\ell-2(\varepsilon_1 w_1 + r\ell))}$ and $W_{-\varepsilon_1, \varepsilon_2}^{(n, m+2\ell-2(\varepsilon_1 w_1 + r\ell))} \rightarrow W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$. Therefore our original pair of cell modules are in the same block.

The proof for $w_2 \in \mathbb{Z}, w_1 \notin \mathbb{Z}$ is similar, except we must consider cases (29) and (33), and use Theorem 8.4 in place of Theorem 8.3. We therefore have the following theorem:

Theorem 9.3. *Suppose q is a 2ℓ -th root of unity and $w_1 \in \mathbb{Z}, w_2 \notin \mathbb{Z}$. Then two cell modules $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ and $W_{\eta_1, \eta_2}^{(n, t)}$ are in the same block if and only if $\varepsilon_2 = \eta_2$ and $|m - \varepsilon_1 w_1 - \varepsilon_2 w_2| \equiv |t - \eta_1 w_1 - \varepsilon_2 w_2| \pmod{2\ell}$.*

If now $w_2 \in \mathbb{Z}, w_1 \notin \mathbb{Z}$, then two cell modules $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ and $W_{\eta_1, \eta_2}^{(n, t)}$ are in the same block if and only if $\varepsilon_1 = \eta_1$ and $|m - \varepsilon_1 w_1 - \varepsilon_2 w_2| \equiv |t - \eta_1 w_1 - \varepsilon_2 w_2| \pmod{2\ell}$.

If q is not a root of unity, then replace the above two congruences modulo 2ℓ by equalities.

Figure 9 shows two plots of the cell modules, when just w_1 and just w_2 are integral respectively. The arrows indicate a homomorphism between the corresponding modules.

9.3. Either $w_1 + w_2$ or $w_1 - w_2$ integral. We first turn to the case $w_1 + w_2 \in \mathbb{Z}$ but $w_1 - w_2 \notin \mathbb{Z}$. Again, we begin by taking q to not be a root of unity. Here, we are looking to satisfy equations

(27) and (31) with $\varepsilon_1 = \varepsilon_2$. Once more, we see that blocks have size at most two. Now if we have a solution to equation (27), then for these modules to be in the same block we must have a non-zero homomorphism

$$W_{+,+}^{(n,m)} \longrightarrow W_{-,-}^{(n,m-2w_1-2w_2)}.$$

However by restricting both modules to the left-blob algebra we see from Table 4 that the two modules have different standard contents. Thus, since $w_1 \notin \mathbb{Z}$, we deduce that there can be no such homomorphism.

In the case of equation (31), we can again assume that $m > t$. Then since m and t are both non-negative integers we can only have a solution if $\varepsilon_1 = \varepsilon_2 = \text{sgn}(w_1 + w_2)$. Therefore we have $m > t = 2(\varepsilon_1 w_1 + \varepsilon_2 w_2) - m \geq 0$ (with equality only if $\varepsilon_1 = \varepsilon_2 = 1$), and thus we can use Theorem 8.5 to show that the cell modules $W_{+,+}^{(n,m)}$ and $W_{+,+}^{(n,t)}$ are in the same block.

If q is a 2ℓ -th root of unity, then we must consider equations (27), (30), (31) and (34). Again, since neither w_1 nor w_2 are integers we can rule out (27) and (34) by restricting to either the left- or right-blob algebra. In the case of (31), we can show that there exists a non-zero homomorphism between the cell modules in the same way as above. It remains, therefore, to deal with (30). We will begin by showing that $W_{\varepsilon_1, \varepsilon_2}^{(n,m)}$ and $W_{\varepsilon_1, \varepsilon_2}^{(n,m+2\ell)}$ are in the same block, and the general result will follow. Indeed, we choose $r \in \mathbb{Z}$ such that $m + 2\ell > 2(\varepsilon_1 w_1 + \varepsilon_2 w_2 + r\ell) - (m + 2\ell) > m$, then by Theorem 8.5 we have non-zero homomorphisms $W_{\varepsilon_1, \varepsilon_2}^{(n,m+2\ell)} \longrightarrow W_{\varepsilon_1, \varepsilon_2}^{(n, 2(\varepsilon_1 w_1 + \varepsilon_2 w_2 + r\ell) - (m + 2\ell))}$ and $W_{\varepsilon_1, \varepsilon_2}^{(n, 2(\varepsilon_1 w_1 + \varepsilon_2 w_2 + r\ell) - (m + 2\ell))} \longrightarrow W_{\varepsilon_1, \varepsilon_2}^{(n,m)}$. Therefore our original pair of cell modules are in the same block.

Theorem 9.4. *Suppose q is a 2ℓ -th root of unity and $w_1 + w_2 \in \mathbb{Z}, w_1 - w_2 \notin \mathbb{Z}$. Then two cell modules $W_{\varepsilon_1, \varepsilon_2}^{(n,m)}$ and $W_{\eta_1, \eta_2}^{(n,t)}$ are in the same block if and only if $\varepsilon_1 = \eta_1 = \varepsilon_2 = \eta_2$ and $|m - \varepsilon_1 w_1 - \varepsilon_2 w_2| \equiv |t - \eta_1 w_1 - \eta_2 w_2| \pmod{2\ell}$.*

If q is not a root of unity then replace the above congruence modulo 2ℓ by an equality.

The case $w_1 - w_2 \in \mathbb{Z}_{>0}$ but $w_1, w_2, w_1 + w_2 \notin \mathbb{Z}$ is proved similarly.

Theorem 9.5. *Suppose q is a 2ℓ -th root of unity and $w_1 - w_2 \in \mathbb{Z}, w_1 + w_2 \notin \mathbb{Z}$. Then two cell modules $W_{\varepsilon_1, \varepsilon_2}^{(n,m)}$ and $W_{\eta_1, \eta_2}^{(n,t)}$ are in the same block if and only if $\varepsilon_1 = \eta_1 = -\varepsilon_2 = -\eta_2$ and $|m - \varepsilon_1 w_1 - \varepsilon_2 w_2| \equiv |t - \eta_1 w_1 - \eta_2 w_2| \pmod{2\ell}$.*

If q is not a root of unity then replace the above congruence modulo 2ℓ by an equality.

Figure 10 shows two plots of the cell modules, when just $w_1 + w_1$ and just $w_1 - w_2$ are integral respectively. The arrows indicate a homomorphism between the corresponding modules.

9.4. Both $w_1 + w_2$ and $w_1 - w_2$ integral. If now we have $w_1 + w_2, w_1 - w_2 \in \mathbb{Z}$ but $w_1, w_2 \notin \mathbb{Z}$, then we must have $w_1, w_2 \in \frac{1}{2}\mathbb{Z} \setminus \mathbb{Z}$. The labels m and t for cell modules all have the same parity, in particular $m \pm t$ is always even, whereas both $2w_1$ and $2w_2$ are odd. Therefore there can be no solutions to (28),(29),(32) nor (33). Thus we simply combine Theorems 9.4 and 9.5.




FIGURE 10. Graphical depiction of the cell modules of (i) b'_9 with $w_1 = \frac{1}{4}$ and $w_2 = \frac{11}{4}$; and (ii) b'_8 with $w_1 = \frac{1}{4}$ and $w_2 = -\frac{7}{4}$.




FIGURE 11. Graphical depiction of the cell modules of b'_8 with $w_1 = \frac{5}{2}$ and $w_2 = -\frac{1}{2}$.

Theorem 9.6. *Suppose q is a 2ℓ -th root of unity and $w_1 + w_2 \in \mathbb{Z}$, $w_1 - w_2 \in \mathbb{Z}$ but $w_1, w_2 \notin \mathbb{Z}$. Then two cell modules $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ and $W_{\eta_1, \eta_2}^{(n, t)}$ are in the same block if and only if $|m - \varepsilon_1 w_1 - \varepsilon_2 w_2| \equiv |t - \eta_1 w_1 - \eta_2 w_2| \pmod{2\ell}$ and either*

- (1) $\varepsilon_1 = \eta_1 = \varepsilon_2 = \eta_2$, or
- (2) $\varepsilon_1 = \eta_1 = -\varepsilon_2 = -\eta_2$.

If q is not a root of unity then replace the above congruence modulo 2ℓ by an equality.

Figure 11 shows a plot the cell modules when both $w_1 + w_2$ and $w_1 - w_2$ are integral, but not w_1 nor w_2 . The arrows indicate a homomorphism between the corresponding modules.

9.5. Both w_1 and w_2 integral. Finally, we consider the case when $w_1, w_2 \in \mathbb{Z}$ and q not a root of unity. As explained in the beginning of this section, we can globalise appropriately so that we only

consider cell modules $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ with $m \ll N$ and $w_1, w_2 > 0$. We may also assume that $w_1 \leq w_2$, since we can swap blobs and flip diagrams horizontally to make this the case. By fixing ε_1 and ε_2 and considering the equations (27)–(34) we see that each block has size at most 4 as q is not a root of unity. By repeated globalisation we will determine the blocks containing modules of the form $W_{+,+}^{(N, m)}$ where $N \gg m$. By then repeated localising, we see that we will in fact deal with each cell module in b'_n . Care must be taken when localising, as we will encounter blocks of size 4 in the large N limit which may break up into singleton blocks when localising back to n .

Consider first the case $m < w_1 + w_2$. Then we see by Theorem 8.5 that we have a homomorphism

$$W_{+,+}^{(N, 2w_1+2w_2-m)} \longrightarrow W_{+,+}^{(N, m)}.$$

Now we are assuming that $0 < w_1 \leq w_2$. Thus $2w_1 + 2w_2 - m > w_1 + w_2 \geq 2w_1$, therefore $2w_1 + 2w_2 - m - 2w_1 = 2w_2 - m > 0$ and we can use Theorem 8.3 to obtain a homomorphism

$$W_{+,+}^{(N, 2w_1+2w_2-m)} \longrightarrow W_{-,+}^{(N, 2w_2-m)}.$$

When localising it is possible that the module $W_{+,+}^{(N, 2w_1+2w_2-m)}$ will be annihilated, in which case the remaining two modules may not be linked via homomorphisms. We now show that this is the case:

Lemma 9.7. *Suppose $q \in \mathbb{C}^\times$ is not a root of unity and $w_1, w_2 \in \mathbb{Z}_{>0}$ be as above. Let $m < w_1 + w_2$ be a non-negative integer. Then we have*

$$\text{Hom}(W_{+,+}^{(n, m)}, W_{-,+}^{(n, 2w_2-m)}) = \text{Hom}(W_{-,+}^{(n, 2w_2-m)}, W_{+,+}^{(n, m)}) = 0.$$

Proof. If we have a non-zero homomorphism between the cell modules, then this must restrict to a homomorphism between cell modules for the left blob algebra. We will show that there can be no such homomorphism.

Suppose first that we have a non-zero homomorphism $W_{+,+}^{(n, m)} \longrightarrow W_{-,+}^{(n, 2w_2-m)}$. By applying the localisation functor F an even number of times (so that no parameter change occurs), we may assume that $n = m + 1$. Now restricting to the left blob algebra, we see from Table 4 that $W_{+,+}^{(m+1, m)}$ has standard content $W_{m+1}(m+1)$ and $W_{-,+}^{(m+1, 2w_2-m)}$ has standard content $W_{-(m+1)}(m+1), W_{-(m-1)}(m+1), \dots, W_{-(2w_2-m+1)}(m+1)$. As such, we must have a left blob homomorphism from the trivial module to one of the latter standard modules. However since q is not a root of unity, the module $W_{m+1}(m+1)$ is mapped only to $W_{2w_1-m-1}(m+1)$. We then note that $2w_1 - m - 1 > w_1 - w_2 - 1$, whereas $-(2w_2 - m + 1) < w_1 - w_2 - 1$. Therefore there is no left-blob homomorphism in the restriction, and thus no symplectic blob homomorphism.

If we now assume that we have a homomorphism $W_{-,+}^{(n, 2w_2-m)} \longrightarrow W_{+,+}^{(n, m)}$, then by applying the localisation functor F an odd number of times we may assume that $n = 2w_2 - m$. Then by restricting to the left blob algebra we see that $W_{+,+}^{(2w_2-m, 2w_2-m-1)}$ has standard content $W_{2w_2-m}(2w_2-m)$, whereas $W_{-,+}^{(2w_2-m, m+1)}$ has standard content $W_{-(2w_2-m)}(2w_2-m), \dots, W_{-(m+2)}(2w_2-m)$. Again

by considering left blob homomorphisms with q not a root of unity, and taking the parameter change $w_1 \mapsto -w_1 - 1$ into account, we see that the trivial module is mapped only to $2(-w_1 - 1) - 2w_2 + m = -2(w_1 + 2w_2) + m - 2$. Note now that this is less than $-2w_2 + m$, which is the least label in the latter set of standard contents. Thus there can be no left blob homomorphism, and hence no symplectic blob homomorphism. $\square$

So far we have found three distinct modules with the same eigenvalue, all linked via homomorphisms.

If also $2w_1 + 2w_2 - m > 2w_2$ (so that $m < 2w_1$), then we can use Theorem 8.4 to obtain a homomorphism

$$W_{+,+}^{(N, 2w_1+2w_2-m)} \longrightarrow W_{+,-}^{(N, 2w_1-m)}.$$

In a manner similar to Lemma 9.7, we can show that there are no homomorphisms between $W_{+,-}^{(N, 2w_1-m)}$ and $W_{-,+}^{(N, 2w_2-m)}$ or $W_{+,+}^{(N, m)}$. Since blocks have size at most four, this block has the structure as in Figure 12, where arrows indicate the existence of a homomorphism and lack of arrows indicates the non-existence of a homomorphism.




FIGURE 12. The block structure for $m < 2w_1$.


Note that if $n \leq 2w_1 + 2w_2 - m$, then the module $W_{+,+}^{(n, 2w_1+2w_2-m)}$ does not exist. Since there are no homomorphisms between the remaining three modules, the block breaks up into singleton blocks.

If now $2w_1 + 2w_2 - m = 2w_2$ (so that $m = 2w_1$), then we claim that there are at most three possible solutions to equations (27)–(34). In particular we can satisfy neither (28) nor (32) and so the block has size at most three, and we have found enough homomorphisms. Since we have fixed $\varepsilon_1 = 1$, equations (28) and (32) reduce to $m \pm t = m$, so that $t = 0$. However the only module that exists when $t = 0$ also must have $\eta_1 = \eta_2 = 1$, which is not valid when considering this pair of equations. We thus have the block structure as in Figure 13, with the same convention for arrows (or lack thereof).

Again we see that if $n \leq 2w_2$ then the module $W_{+,+}^{(n, 2w_2)}$ does not exist. Therefore the block once more breaks up into singleton blocks.

Finally, if $2w_1 + 2w_2 - m < 2w_2$, then we have $m > 2w_1$ and so we can again use Theorem 8.3 to obtain


$$W_{+,+}^{(N, m)} \longrightarrow W_{-,+}^{(N, m-2w_1)}.$$

FIGURE 13. The block structure for $m = 2w_1$.

We also see that the modules $W_{-,+}^{(N, 2w_2-m)}$ and $W_{-,+}^{(N, m-2w_1)}$ satisfy the conditions of Theorem 8.5, and thus have a homomorphism

$$W_{-,+}^{(N, 2w_2-m)} \longrightarrow W_{-,+}^{(N, m-2w_1)}.$$

Again, since blocks have size at most four, this block has the structure as in Figure 14, where again the arrows indicate the existence of a homomorphism, the lack of arrows the non-existence, and a dotted line indicates an unknown (which will not matter when considering the block structure).

FIGURE 14. The block structure for $2w_1 < m < w_1 + w_2$

As in the previous two cases, if $n \leq 2w_1 + 2w_2 - m$ then the module $W_{+,+}^{(n, 2w_1+2w_2-m)}$ no longer exists. However since we have homomorphisms between the remaining modules, the block does not decompose further. The same is true if $n \leq m$ or $n < 2w_2 - m$.

We have now accounted for all modules $W_{+,+}^{(n, m)}$ with $0 \leq m \leq 2w_1 + 2w_2$ ($m \neq w_1 + w_2$), $W_{-,+}^{(n, m)}$ with $0 < m \leq 2w_2$ ($m \neq w_2 - w_1$), and $W_{+,+}^{(n, m)}$ with $0 < m \leq 2w_1$.

Suppose now $m = w_1 + w_2$, again with $w_1 \leq w_2$. In this case, we show that there are at most two modules in the block and find a homomorphism between them. Indeed, from equations (27)–(34) the only non-trivial solution is $l = w_2 - w_1$, $\varepsilon_1 \neq \eta_1$, $\varepsilon_2 = \eta_2$. But then this satisfies the conditions of Theorem 8.3, and we have a homomorphism

$$W_{+,+}^{(N, w_1+w_2)} \longrightarrow W_{-,+}^{(N, w_2-w_1)}.$$

This gives a block structure as in Figure 15.

This means that now we have covered all modules $W_{+,+}^{(n, m)}$ with $0 \leq m \leq 2w_1 + 2w_2$, $W_{-,+}^{(n, m)}$ with $0 < m \leq 2w_2$, and $W_{+,+}^{(n, m)}$ with $0 < m \leq 2w_1$.

$$\begin{array}{c}
W_{+,+}^{(N,w_1+w_2)} \\
\downarrow \\
W_{-,+}^{(N,w_2-w_1)}
\end{array}$$

FIGURE 15. The block structure for $m = w_1 + w_2$.

So it remains to consider the case $m > 2w_1 + 2w_2$. Here we have both $2w_1 < m$ and $2w_2 < m$, so both Theorems 8.3 and 8.4 are satisfied and we have homomorphisms

$$\begin{aligned}
W_{+,+}^{(N,m)} &\longrightarrow W_{-,+}^{(N,m-2w_1)}, \text{ and} \\
W_{+,+}^{(N,m)} &\longrightarrow W_{+,-}^{(N,m-2w_2)}.
\end{aligned}$$

But in this case we also have $m - 2w_2 > 2w_1$ and $m - 2w_1 > 2w_2$, so we can again use Theorems 8.3 and 8.4 to obtain homomorphisms

$$\begin{aligned}
W_{+,-}^{(N,m-2w_2)} &\longrightarrow W_{-,-}^{(N,m-2w_1-2w_2)}, \text{ and} \\
W_{-,+}^{(N,m-2w_1)} &\longrightarrow W_{-,-}^{(N,m-2w_1-2w_2)}.
\end{aligned}$$

Note that this final case deals with all modules $W_{+,+}^{(n,m)}$ with $m > 2w_1 + 2w_2$, $W_{-,+}^{(n,m)}$ with $m > 2w_2$, $W_{+,-}^{(m,n)}$ with $m > 2w_1$ and $W_{-,-}^{(n,m)}$ with $m > 0$. In this case, the block structure is shown in Figure 16.

$$\begin{array}{ccccc}
& & W_{+,+}^{(N,m)} & & \\
& \swarrow & & \searrow & \\
W_{-,+}^{(N,m-2w_1)} & \cdots & & \cdots & W_{+,-}^{(N,m-2w_2)} \\
& \swarrow & & \searrow & \\
& & W_{-,-}^{(N,m-2w_1-2w_2)} & &
\end{array}$$

FIGURE 16. The block structure for large m .

The cases displayed in Figures 12–16 above exhaust the list of modules and we are able to give the final main result.

Theorem 9.8. *Suppose q is not a root of unity and $w_1, w_2 \in \mathbb{Z}$. Let $\sigma_1 = \text{sgn}(w_1)$, $\sigma_2 = \text{sgn}(w_2)$. Then for $n \geq 2|w_1| + 2|w_2| + \frac{1}{2}(\sigma_1 + \sigma_2)$, two cell modules $W_{\varepsilon_1, \varepsilon_2}^{(n,m)}$ and $W_{\eta_1, \eta_2}^{(n,l)}$ are in the same block if and only if $|m - \varepsilon_1 w_1 - \varepsilon_2 w_2| = |l - \eta_1 w_1 - \eta_2 w_2|$.*

Proof. In the case $n \geq 2|w_1| + 2|w_2| + \frac{1}{2}(\sigma_1 + \sigma_2)$, none of the blocks described by Figures 12 and 13 break up into singleton blocks. Therefore in the globalised case with $w_1, w_2 > 0$, the Theorem follows. Notice that

$$\begin{aligned} |m - \varepsilon_1 w_1 - \varepsilon_2 w_2| &= |(m + \varepsilon_1) - (-\varepsilon_1)(-w_1 - 1) - \varepsilon_2 w_2| \\ &= |(m + \varepsilon_2) - \varepsilon_1 w_1 - (-\varepsilon_2)(-w_2 - 1)| \\ &= |(m + \varepsilon_1 + \varepsilon_2) - (-\varepsilon_1)(-w_1 - 1) - (-\varepsilon_2)(-w_2 - 1)|, \end{aligned}$$

and so the result holds after localising. $\square$

In the case $n < 2|w_1| + 2|w_2| + \frac{1}{2}(\sigma_1 + \sigma_2)$, we do not have a succinct characterisation of the blocks of b'_n . We therefore make the following statement.

Theorem 9.9. *Suppose q is not a root of unity and $w_1, w_2 \in \mathbb{Z}$. Let $\sigma_1 = \text{sgn}(w_1), \sigma_2 = \text{sgn}(w_2)$. Then for $n < 2|w_1| + 2|w_2| + \frac{1}{2}(\sigma_1 + \sigma_2)$, the blocks of b'_n are obtained by considering the blocks of the algebra b'_N with $N \gg n$, $w_1, w_2 > 0$ as in Figures 12–16, then localising back to b'_n . The blocks are then given by removing any annihilated modules and associated arrows from the appropriate figure and taking the connected components of what remains.*

Turning now to the case when q is a 2ℓ -th root of unity, we will determine blocks in the large N limit. Here, we will show that any pair of cell modules satisfying equations (27)–(34) are in the same block. Therefore in what follows, assume that $W_{\varepsilon_1, \varepsilon_2}^{(N, m)}$ and $W_{\eta_1, \eta_2}^{(N, t)}$ are two cell modules satisfying the current equation in question. This will determine η_1, η_2 in terms of $\varepsilon_1, \varepsilon_2$, and also the congruence class of t modulo 2ℓ .

We deal with equations (28), (29), (30) and (31) simultaneously. By applying the arguments from Sections 9.2 and 9.3, we see that $W_{\varepsilon_1, \varepsilon_2}^{(N, m)}$ and $W_{\eta_1, \eta_2}^{(N, t)}$ are in the same block.

Now consider equation (27). In this case we choose $r \in \mathbb{Z}$ such that $m - 2(\varepsilon_1 w_1 + r\ell) > 0$ and consider the cell module $W_{-\varepsilon_1, \varepsilon_2}^{(N, m - 2(\varepsilon_1 w_1 + r\ell))}$. We then see that $W_{-\varepsilon_1, \varepsilon_2}^{(N, m - 2(\varepsilon_1 w_1 + r\ell))}$ and $W_{\varepsilon_1, \varepsilon_2}^{(N, m)}$ satisfy (28), so are in the same block by previous arguments. Moreover $W_{-\varepsilon_1, \varepsilon_2}^{(N, m - 2(\varepsilon_1 w_1 + r\ell))}$ and $W_{\eta_1, \eta_2}^{(N, t)}$ satisfy (29) (due to our assumptions on η_1, η_2 and t), and are therefore in the same block. By transitivity, we see that the original two modules are in the same block.

If the two modules satisfy (32), then then we will again link these modules via a third. In particular we consider $W_{\varepsilon_1, \varepsilon_2}^{(N, 2(\varepsilon_1 w_1 + \varepsilon_2 w_2 + r\ell) - m)}$, where $r \in \mathbb{Z}$ is chosen so that $2(\varepsilon_1 w_1 + \varepsilon_2 w_2 + r\ell) - m > 0$. Then this module and $W_{\varepsilon_1, \varepsilon_2}^{(N, m)}$ satisfy (31) and are therefore in the same block. Moreover, $W_{\varepsilon_1, \varepsilon_2}^{(N, 2(\varepsilon_1 w_1 + \varepsilon_2 w_2 + r\ell) - m)}$ and $W_{\eta_1, \eta_2}^{(N, t)}$ satisfy (29) and are in the same block. Therefore the original pair of modules are in the same block.

If the two modules satisfy (33) then we use an argument analogous to the previous paragraph.

Finally, if two modules satisfy (34), then we again use a third module to show that the original two are in the same block. In particular we choose $r \in \mathbb{Z}$ so that $m - 2(\varepsilon_1 w_1 + r\ell) > 0$, then




FIGURE 17. Graphical depiction of the cell modules of b'_{13} with $w_1 = 3$ and $w_2 = 1$.

we see that $W_{\varepsilon_1, \varepsilon_2}^{(N, m)}$ and $W_{-\varepsilon_1, -\varepsilon_2}^{(N, m-2(\varepsilon_1 w_1 + r\ell))}$ are in the same block (as they satisfy (28)), and also $W_{-\varepsilon_1, -\varepsilon_2}^{(N, m-2(\varepsilon_1 w_1 + r\ell))}$ and $W_{\eta_1, \eta_2}^{(N, t)}$ are in the same block (as they satisfy (32)).

We therefore arrive at the following theorem:

Theorem 9.10. *Suppose q is a 2ℓ -th root of unity and $w_1, w_2 \in \mathbb{Z}$. Then for $N \gg \max\{m, t\}$, two cell modules $W_{\varepsilon_1, \varepsilon_2}^{(N, m)}$ and $W_{\eta_1, \eta_2}^{(N, t)}$ are in the same block if and only if $|m - \varepsilon_1 w_1 - \varepsilon_2 w_2| \equiv |t - \eta_1 w_1 - \eta_2 w_2| \pmod{2\ell}$.*

Figure 17 shows a plots of the cell modules when both w_1 and w_2 are integral. The arrows indicate a homomorphism between the corresponding modules, and the dashed square indicate that these modules are in the same block. Note that away from the extremes of each arm, there is a uniform pattern of concentric squares.

10. LINKAGE VIA THE MODULE $W^{(n)}(b)$

In the above we have been working with cell modules for $b'_n := b_n^x / I_n(0)$. This precisely excludes the 2^n -dimensional module $W^{(n)}(b)$. We will now deal with linkage via this module in b_n^x and thus complete our investigation into the block structure. We begin with the following theorem:

Theorem 10.1 ([5, Theorem 5.17]). *The Gram determinant Γ_b^n of $W^{(n)}(b)$ (with respect to the path basis, as defined in (25)) is given by:*

- for n even:

$$\Gamma_b^n = \alpha_n \prod_{m=0}^{(n-2)/2} \left(\prod_{\varepsilon_1, \varepsilon_2, \varepsilon_3 = \pm 1} [(1 + 2m + \varepsilon_1 w_1 + \varepsilon_2 w_2 + \varepsilon_3 \theta)/2] \right)^{\sum_{i=1}^{(n-2m)/2} \binom{n}{(n-2m-2i)/2}};$$

- for n odd:

$$\begin{aligned} \Gamma_b^n = \alpha_n & \left(\prod_{\varepsilon_2, \varepsilon_3 = \pm 1} [(w_1 + \varepsilon_2 w_2 + \varepsilon_3 \theta)/2] \right)^{2^{n-1}} \\ & \times \prod_{m=1}^{(n-1)/2} \left(\prod_{\varepsilon_1, \varepsilon_2, \varepsilon_3 = \pm 1} [(2m + \varepsilon_1 w_1 + \varepsilon_2 w_2 + \varepsilon_3 \theta)/2] \right)^{\sum_{i=1}^{(n-2m+1)/2} \binom{n}{(n-2m-2i+1)/2}}, \end{aligned}$$

where α_n is given in both cases by

$$\alpha_n = ([w_1][w_2 + 1])^{-2 \sum_{m=0}^{n-1} (\sum_{i=1}^{m+1} \binom{n}{m-i+1})},$$

up to factors that are units under our standing assumptions.

Assuming $\alpha_n \neq 0$, we therefore see that the module $W^{(n)}(b)$ is irreducible unless θ is congruent to $\pm(-m + \varepsilon_1 w_1 + \varepsilon_2 w_2)$ modulo 2ℓ for some integer m . Since this module has label larger than all the other cell modules in the poset ordering, being irreducible implies that it is alone in its block. (This remains true even in the non-quasi-hereditary case as this module has dimension larger than all the other cell modules.) We will say that if $\theta \equiv \pm(-m + \varepsilon_1 w_1 + \varepsilon_2 w_2) \pmod{2\ell}$ then θ is *critical*. If θ is not critical, then the module $W^{(n)}(b)$ is always a singleton block and the algebra b_n^x has the same blocks as b'_n with the added singleton block. Therefore we henceforth suppose that θ is critical.

Suppose that we have two non-isomorphic submodules $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$, $W_{\eta_1, \eta_2}^{(n, t)}$ of $W^{(n)}(b)$. These each correspond to a singular factor of the gram determinant, Γ_b^n . The condition for two of the factors in the gram determinant to be equal to zero is the same condition for the central element Z_n to act by the same constant on the modules $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ and $W_{\eta_1, \eta_2}^{(n, t)}$ with the added condition that $[(m - \varepsilon_1 w_1 - \varepsilon_2 w_2 + \theta)/2] = 0$. This is not so surprising as if the modules $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$, $W_{\eta_1, \eta_2}^{(n, t)}$ appear in the indecomposable module $W^{(n)}(b)$ they must be in the same block. This does mean, that a careful rereading of the proofs of the blocks for the algebra b'_n will give the blocks for the full algebra b_n^x .

We have several cases. In each case, the blocks for b_n^x are the same as for the algebra b'_n , except for the block that contains $W^{(n)}(b)$. In most cases, the module now joins two blocks from b'_n together, except for the case where the modules that have the same action by the central element, already form a block.

To state our result we will only specify the block that can change, i.e. the one that contains the module $W^{(n)}(b)$. The blocks that don't contain $W^{(n)}(b)$ remain the same as for b'_n .

Theorem 10.2. *The blocks of the symplectic blob algebra, b_n^x , when θ has a critical value, $\theta = -m + \epsilon_1 w_1 + \epsilon_2 w_2$, are as follows:*

- (i) *If q is not a root of unity and none of $w_1, w_2, w_1 \pm w_2$ are integral, then the only non-singleton block is the one containing $W^{(n)}(b)$ and then $W^{(b)}(n)$ and $W_{\epsilon_1, \epsilon_2}^{(n, m)}$ are in the same block.*
- (ii) *If q is a primitive 2ℓ -th root of unity and none of $w_1, w_2, w_1 \pm w_2$ are integral, then the module $W^{(n)}(b)$ is in the same block as all $W_{\eta_1, \eta_2}^{(n, t)}$ with $\epsilon_1 = \eta_1, \epsilon_2 = \eta_2$ and $m \equiv t \pmod{2\ell}$ and all $W_{\eta'_1, \eta'_2}^{(n, t')}$ with $\epsilon_1 = -\eta'_1, \epsilon_2 = -\eta'_2$ and $m \equiv -t' \pmod{2\ell}$.*
- (iii) *If $w_1 + w_2 \in \mathbb{Z}$ and none of $w_1, w_2, w_1 - w_2$ are integral, then the module $W^{(n)}(b)$ is in the same block as all $W_{\eta_1, \eta_2}^{(n, t)}$ with $\epsilon_1 = \eta_1 = \epsilon_2 = \eta_2$ and $|m - \epsilon_1 w_1 - \epsilon_2 w_2| \equiv |t - \eta_1 w_1 - \eta_2 w_2| \pmod{2\ell}$, and all $W_{\eta'_1, \eta'_2}^{(n, t')}$ with $\epsilon_1 = -\eta'_1 = \epsilon_2 = -\eta'_2$ and $|m - \epsilon_1 w_1 - \epsilon_2 w_2| \equiv |t - \eta'_1 w_1 - \eta'_2 w_2| \pmod{2\ell}$, where the q is a primitive 2ℓ -th root of unity, or the congruence is an equality if q is not an 2ℓ -th root of unity.*
- (iv) *If $w_1 - w_2 \in \mathbb{Z}$ and none of $w_1, w_2, w_1 + w_2$ are integral, then the module $W^{(n)}(b)$ is in the same block as all $W_{\eta_1, \eta_2}^{(n, t)}$ with $\epsilon_1 = \eta_1 = -\epsilon_2 = -\eta_2$ and $|m - \epsilon_1 w_1 - \epsilon_2 w_2| \equiv |t - \eta_1 w_1 - \eta_2 w_2| \pmod{2\ell}$, and all $W_{\eta'_1, \eta'_2}^{(n, t')}$ with $\epsilon_1 = -\eta'_1 = -\epsilon_2 = \eta'_2$ and $|m - \epsilon_1 w_1 - \epsilon_2 w_2| \equiv |t - \eta'_1 w_1 - \eta'_2 w_2| \pmod{2\ell}$, where the q is a primitive 2ℓ -th root of unity, or the congruence is an equality if q is not an 2ℓ -th root of unity.*
- (v) *If $w_1 + w_2$ and $w_1 - w_2 \in \mathbb{Z}$ and none of w_1, w_2 , are integral, then the module $W^{(n)}(b)$ is in the same block as all $W_{\eta_1, \eta_2}^{(n, t)}$ with $|m - \epsilon_1 w_1 - \epsilon_2 w_2| \equiv |t - \eta_1 w_1 - \eta_2 w_2| \pmod{2\ell}$, where the q is a primitive 2ℓ -th root of unity, or the congruence is an equality if q is not an 2ℓ -th root of unity.*
- (vi) *If $w_1 \in \mathbb{Z}$ or $w_2 \in \mathbb{Z}$ (but not both) and none of $w_2, w_1 \pm w_2$ are integral, then the module $W^{(n)}(b)$ is in the same block as all $W_{\eta_1, \eta_2}^{(n, t)}$ with $|m - \epsilon_1 w_1 - \epsilon_2 w_2| \equiv |t - \eta_1 w_1 - \eta_2 w_2| \pmod{2\ell}$, where the q is a primitive 2ℓ -th root of unity, or the congruence is an equality if q is not an 2ℓ -th root of unity.*
- (vii) *If both $w_1, w_2 \in \mathbb{Z}$ then the module $W^{(n)}(b)$ is in the same block as all $W_{\eta_1, \eta_2}^{(n, t)}$ with $|m - \epsilon_1 w_1 - \epsilon_2 w_2| \equiv |t - \eta_1 w_1 - \eta_2 w_2| \pmod{2\ell}$, where the q is a primitive 2ℓ -th root of unity, or the congruence is an equality if q is not an 2ℓ -th root of unity. (This is the only case where the module $W^{(n)}(b)$ does not join two blocks from b'_n .)*

Proof. (i). Clear as all $-m + \epsilon_1 w_1 + \epsilon_2 w_2 - \theta$ are distinct in this case and considering the Gram determinant of $W^{(n)}(b)$.

(ii). The module $W_{\epsilon_1, \epsilon_2}^{(n, m)}$ embeds into $W^{(n)}(b)$, by the assumption on θ . Thus $W^{(n)}(b)$ is in the same block as all $W_{\eta_1, \eta_2}^{(n, t)}$ that are linked to $W_{\epsilon_1, \epsilon_2}^{(n, m)}$. Pick $a \in \mathbb{Z}$ such that $0 \leq 2a\ell - m \leq n$. Then $[(2a\ell - m + \epsilon_1 w_1 + \epsilon_2 w_2 - \theta)/2] = 0$, so $W_{-\epsilon_1, -\epsilon_2}^{(n, 2a\ell - m)}$ is a submodule of $W^{(n)}(b)$. Hence $W^{(n)}(b)$ is in the same block as all $W_{\eta'_1, \eta'_2}^{(n, t')}$ that are linked to $W_{-\epsilon_1, -\epsilon_2}^{(n, 2a\ell - m)}$.

Thus using the proof of Theorem 9.2, the block containing $W^{(n)}(b)$ is the same as the solutions to equations (30) and (34).

(iii). As in (ii), $W^{(n)}(b)$ is in the same block as $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$ and this gives the first condition. Pick $a \in \mathbb{Z}$ such that $0 \leq 2a\ell + m - 2\varepsilon_1(w_1 + w_2) \leq n$. (NB: ℓ is taken to be zero if q is not a primitive 2ℓ -th root of unity. Of course, then such an a may not exist, but then there are no further solutions to the equations that need to be considered.) Consider $t = m - 2\varepsilon_1(w_1 + w_2)$ and the module $W_{-\varepsilon_1, -\varepsilon_1}^{(n, t)}$. For this module $[(t + \varepsilon_1 w_1 + \varepsilon_2 w_2 + \theta)/2] = 0$, so $W_{-\varepsilon_1, -\varepsilon_1}^{(n, t)}$ is a submodule of $W^{(n)}(b)$ and thus they are in the same block. This gives the second condition using Theorem 9.4. These two conditions combined give all solutions to equations (27), (30), (31) and (34), thus this is the whole block.

(iv). This is similar to (iii).

(v). This merges (iii) and (iv) and says that the block is determined by the action of the central element.

(vi). As in (ii), $W^{(n)}(b)$ is in the same block as $W_{\varepsilon_1, \varepsilon_2}^{(n, m)}$. Pick $a \in \mathbb{Z}$ such that $0 \leq 2a\ell - m + 2\varepsilon_1 w_1 \leq n$. (NB: ℓ is taken to be zero if q is not a primitive 2ℓ -th root of unity. Of course, then such an a may not exist, but then there are no further solutions to the equations that need to be considered.) Consider $t = -m + 2\varepsilon_1 w_1$ and the module $W_{\varepsilon_1, -\varepsilon_2}^{(n, t)}$. For this module $[(-t + \varepsilon_1 w_1 - \varepsilon_2 w_2 + \theta)/2] = 0$, so $W_{\varepsilon_1, -\varepsilon_2}^{(n, t)}$ is a submodule of $W^{(n)}(b)$ and thus they are in the same block. This gives the condition as stated using Theorem 9.3, which is the same as the condition for the central element to act by zero.

(vii). Clear as the block is already determined by the action of the central element. $\square$

APPENDIX A. REDUCTION TO HOM SPACES

It is well known that we may identify the blocks of a finite dimensional algebra with the connected components of the Ext^1 quiver between simple modules. Here we prove that finding these connected components is equivalent to determining the connected components of the Hom quiver between standards. Thus in determining the blocks, it is only necessary to compute enough homomorphisms between standard modules in order to find these connected components.

(Here by the Hom quiver between standards, we mean take the quiver whose vertices are labelled by standard modules, $\Delta(\mu)$ and with the number of arrows from λ to μ equal to the dimension of $\text{Hom}(\Delta(\lambda), \Delta(\mu))$.)

Proposition A.1. *Let A be a quasi-hereditary algebra with a simple preserving duality and poset $(\Lambda, \leq)$. For $\lambda \in \Lambda$, let the standard modules be denoted by $\Delta(\lambda)$, costandards by $\nabla(\lambda)$, the principal indecomposable modules by $P(\lambda)$, the irreducible head of this module by $L(\lambda)$ and the indecomposable injective hulls by $I(\lambda)$.*

The blocks of A may be identified with the connected components of the Hom quiver between standards.

Proof. Now if $\text{Hom}(\Delta(\lambda), \Delta(\mu))$ is non-zero then $L(\lambda)$ must be a composition factor of $\Delta(\mu)$ as $\Delta(\lambda)$ has simple head $L(\lambda)$. Thus $L(\lambda)$ and $L(\mu)$ are in the same block. Thus the connected components of the Ext^1 quiver on simples are disjoint unions of the connected components of the Hom quiver on standards.

We now prove the converse, that the connected components of the Hom quiver on standards are disjoint unions of the Ext^1 quiver on simples. Let λ and $\mu \in \Lambda$ with $\text{Ext}^1(L(\lambda), L(\mu))$. Without loss of generality we may assume that $\lambda \geq \mu$ as A has a simple preserving duality (and hence $\text{Ext}^1(L(\lambda), L(\mu)) = \text{Ext}^1(L(\mu), L(\lambda))$). Since the extension of $L(\lambda)$ by $L(\mu)$ has simple socle and $\lambda \geq \mu$, this extension must be a quotient of $\Delta(\lambda)$ and in particular $[\Delta(\lambda) : L(\mu)] \neq 0$. Thus $\text{Hom}(P(\mu), \Delta(\lambda))$ is non-zero, as the dimension of $\text{Hom}(P(\mu), \Delta(\lambda))$ is equal to $[\Delta(\lambda) : L(\mu)]$.

Now if $\text{Hom}(\Delta(\mu), \Delta(\lambda))$ is non-zero then we are done, so suppose that $\text{Hom}(\Delta(\mu), \Delta(\lambda)) = 0$. As $\text{Hom}(P(\mu), \Delta(\lambda))$ is non-zero, there is a submodule of $\Delta(\lambda)$, M , which is a quotient of $P(\mu)$ and hence has simple head $L(\mu)$. As $\text{Hom}(\Delta(\mu), \Delta(\lambda))$ is zero, this quotient M must contain a composition factor, $L(\nu)$ say, for which $\nu \not\leq \mu$. Let $L(\nu)$ be a largest composition factor in M .

Now as M is a submodule of $\Delta(\lambda)$ we must have that $\nu < \lambda$. (It cannot be equal as then M would be the whole of $\Delta(\lambda)$ contradicting $\text{Hom}(\Delta(\mu), \Delta(\lambda)) = 0$.) Now take the largest submodule of M with $L(\nu)$ as its simple head. As ν was maximal, this submodule has composition factors strictly less than ν and hence is a quotient of $\Delta(\nu)$ and so $\text{Hom}(\Delta(\nu), \Delta(\lambda)) \neq 0$.

Also, as M is not a quotient of $\Delta(\mu)$, it must contain a proper submodule N which is a quotient of the kernel Q , of the projection map from $P(\mu)$ to $\Delta(\mu)$. This proper submodule N must contain the $L(\nu)$ as $L(\nu)$ is not a composition factor of $\Delta(\mu)$. As ν is maximal, this ν must be the head of some Δ appearing in a Δ -filtration of Q . I.e. $\Delta(\nu)$ is a section of Q . Thus $\text{Hom}(P(\mu), \nabla(\nu))$ is nonzero. Using the duality we then have $\text{Hom}(\Delta(\nu), I(\mu)) \neq 0$, which implies that $L(\mu)$ is a composition factor of $\Delta(\mu)$ and thus that $\text{Hom}(P(\mu), \Delta(\nu))$ is non-zero.

If $\text{Hom}(\Delta(\mu), \Delta(\nu))$ is non-zero we may stop, otherwise we repeat the argument until we have a chain of ν_i 's with $\lambda > \nu_1 > \nu_2 > \cdots > \nu_m$ and $\text{Hom}(\Delta(\nu_i), \Delta(\nu_{i+1})) \neq 0$. Since Λ is finite this chain must stop eventually with $\text{Hom}(\Delta(\nu_m), \Delta(\mu)) \neq 0$. We may thus conclude that λ and μ are in the same connected component of the Hom quiver on standards. $\square$

REFERENCES

1. R. J. Baxter, *Exactly solved models in statistical mechanics*, Academic, 1981.
2. D. J. Benson, *Representations and Cohomology I*, Cambridge Studies in Advanced Mathematics, no. 30, Cambridge University Press, 1995.
3. R. Brauer, *On modular and p -adic representations of algebras*, Proceedings of the National Academy of Sciences **25** (1939), no. 5, 252–258.

4. A. G. Cox, J. J. Graham, and P. P. Martin, *The blob algebra in positive characteristic*, J. Algebra **266** (2003), 584–635.
5. J. de Gier and A. Nichols, *The two-boundary Temperley-Lieb algebra*, J. Algebra **321** (2009), no. 4, 1132–1167.
6. J. Graham and G. Lehrer, *Cellular algebras*, Inventiones Math. **123** (1996), 1–34.
7. R. M. Green, P. P. Martin, and A. E. Parker, *Towers of recollement and bases for diagram algebras: planarity and beyond*, J. Algebra **316** (2007), 392–452.
8. ———, *A presentation for the symplectic blob algebra*, J. Algebra Appl. **11** (2012), no. 3, 1250060, 22.
9. ———, *On quasi-heredity and cell module homomorphisms in the symplectic blob algebra*, arXiv (2015).
10. R. Hartshorne, *Algebraic geometry*, Springer, 1977.
11. J. E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge Studies in Advanced Mathematics, no. 30, Cambridge University Press, 1990.
12. J. L. Jacobsen and H. Saleur, *Combinatorial aspects of boundary loop models*, J. Stat. Mech. (2008).
13. J. C. Jantzen, *Representations of Algebraic Groups*, Pure Appl. Math., vol. 131, Academic Press, San Diego, 1987.
14. G. Lusztig, *Left cells in Weyl groups*, LNM 1024, Springer (1983).
15. ———, *Lectures on Hecke algebras with unequal parameters*, MIT Lecture notes (1999).
16. P. P. Martin, *Potts models and related problems in statistical mechanics*, vol. 5, World Scientific, 1991.
17. P. P. Martin and H. Saleur, *The blob algebra and the periodic Temperley-Lieb algebra*, Lett. Math. Phys. **30** (1994), no. 3, 189–206.
18. P. P. Martin and D. Woodcock, *On the structure of the blob algebra*, J. Algebra **225** (2000), 957–988.
19. A. Reeves, *Tilting modules for the symplectic blob algebra*, arXiv:1111.0146 (2012).
20. H. N. V. Temperley and E. H. Lieb, *Relations between the “percolation” and “colouring” problem and other graph-theoretical problems associated with regular planar lattices: some exact results for the “percolation” problem*, Proc. Roy. Soc. London Ser. A **322** (1971), no. 1549, 251–280.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS, LS2 9JT, UK

E-mail address: O.H.King@leeds.ac.uk

E-mail address: ppmartin@maths.leeds.ac.uk

E-mail address: A.E.Parker@leeds.ac.uk